

This file contains:

- 1. Sherwood Lake - Lake Classification Report (2008)**
- 2. Sherwood Lake - Aquatic Plant Report with Management Recommendations (2014)**
- 3. Tri-Lakes Management Plan Table (2015)**

SHERWOOD LAKE

LAKE CLASSIFICATION REPORT



**Presented by Reesa Evans, CLM, Lake Specialist
Adams County Land & Water Conservation Department
P.O. Box 287, Friendship, WI 53934**

NOVEMBER 2008

**SHERWOOD LAKE
LAKE CLASSIFICATION REPORT
TABLE OF CONTENTS**

Executive Summary	1
Recommendations	8
Introduction	10
Methods of Data Collection	11
Water Quality Computer Modeling	12
Dissemination of Project Deliverables	12
Adams County Information	13
Figure 1: Location Map of Adams County	13
Sherwood Lake Information	14
Figure 2: Location Map of Sherwood Lake	14
Figure 3: Archeological Map	15
Figure 4: Sherwood Watershed Soils	17
Prior Studies of Sherwood Lake or Tri-Lakes Area	18

Current Land Use	22
Figure 5: Table of Land Use	22
Figure 6: Prior Land Use Table	23
Figure 7a: Surface Watershed Land Use Map	24
Figure 7b: Ground Watershed Land Use Map	25
Figure 8a: Surface Land Use Graph	26
Figure 8b: Ground Land Use Graph	26
Wetlands	27
Figure 9: Example of Lake Wetland	27
Shorelands	28
Figure 10: Graph of Sherwood Shores	28
Figure 11: Photo of Shore Erosion on Sherwood	28
Figure 12: Shoreline Map of Sherwood Lake	29
Figure 13: Buffer map of Sherwood Lake	31
Figure 14: Example of Inadequate Buffer	32
Figure 15: Example of Adequate Buffer	33
Figure 16: Eroded Shore on Sherwood Lake	33
Water Quality	34
Phosphorus	34
Figure 17a: Graph of Phosphorus 1991-2005	35
Figure 17b: Graph of Phosphorus 2006-2007	35
Figure 18: Average Eplimnetic Phosphorus	36
Figure 19a: Table of Current P Loading	37
Figure 19b: Table of Prior P Loading	37
Figure 19c: Contribution by Land Use	38
Figure 20: Graph of Impact of Dec. or Inc.	39
Figure 21: Graph of In-Lake Impact	39
Figure 22: Photo of Algal Bloom	40
Water Clarity	40
Figure 23a: Secchi Readings 1986-1989	40
Figure 23b: Secchi Readings 1990-1999	41

Figure 23c: Secchi Readings 2002-2007	41
Figure 24: Average Growing Season Secchi	42
Figure 25: Photo of Secchi Testing	42
Chlorophyll-a	43
Figure 26a: Graph of Chlorophyll-a Readings	43
Figure 26b: Chlorophyll-a Readings 2006-2007	44
Figure 27: Average Annual Chlorophyll-a	44
Dissolved Oxygen	45
Figure 28a: DO Graph for 2004	46
Figure 28b: DO Graph for 2005	47
Figure 28c: DO Graph for 2006	48
Figure 29a & b: Fish in Big Roche a Cri Lake	49
Hardness, Alkalinity & pH	50
Figure 30: Table of Hardness Levels	50
Figure 31: Graph of County Hardness	50
Figure 32: Table of Acid Rain Sensitivity	51
Figure 33: Graph of County Alkalinity	52
Figure 34: Table of Effects of pH on Fish	52
Figure 35: Graph of pH by Depth	53
Other Water Quality Testing Results	54
Chloride	54
Figure 36: Chloride Levels Graph	54
Nitrogen	54
Calcium & Magnesium	55
Sodium & Potassium	55
Sulfate	56
Figure 37: Graph of Sulfate Levels	56
Turbidity	56
Figure 38: Graph of Turbidity Levels	57
Figure 39: Examples of Turbid Water	57

Hydrologic Budget	58
Figure 40: Example of Hydrologic Budget	58
Figure 41: Bathymetric Map of Sherwood Lake	59
Trophic State	60
Figure 42: Trophic Status Table	60
Figure 43: Sherwood Trophic Status Table	61
Figure 44: Trophic State Index Graph	62
Figure 45: Current TSI v. 1991 TSI	62
In-Lake Habitat	63
Aquatic Plants	63
Figure 46: Aquatic Plant Types in Sherwood	63
Figure 47: Table of Plants Found in 2006	64
Figure 48: Table of Species Richness	66
Figure 49: Table of AMCI 2000 & 2006	67
Figure 50: Changes in Aquatic Plants	70
Figure 51a: Distribution Map of Emergents	71
Figure 51b: Distribution Map of Submergents	71
Figure 52: Table of Herbicide Applications	72
Figure 53: Table of Mechanical Harvesting	73
Figure 54: Photos of Common Native Plants	74
Aquatic Invasives	74
Figure 55: Distribution Map of Exotics	77
Figure 56: Photos of the Exotics in Sherwood	78
Fishery, Wildlife and Endangered Resources	79
Figure 57: Photos of Some of Species of Concern	80
Resources	81
Figure 58: Watercress on Sherwood Lake	82

EXECUTIVE SUMMARY

Background Information about Sherwood Lake

Sherwood Lake is located in the Town of Rome, Adams County, WI, in the south central part of Wisconsin. The impoundment of 14-Mile Creek is slightly over 243 surface acres in size. Maximum depth is 24', with an average depth of 8'. Both Upper and Lower Camelot Lakes flow through dams into Sherwood Lake. Sherwood Lake flows through a dam into Arrowhead Lake. All the Tri-Lakes dams are owned and operated by Adams County. There is a public boat launch on Sherwood Lake on the southwest edge of the lake owned by the Parks Department of Adams County. Heavy residential development around the lake is found along most of the lakeshore. Sherwood Lake is managed by the Tri-Lakes Management District. There is also an active Sherwood Property Owners Association.

The primary soil type in both the surface and ground watersheds is loamy sand. The second most common soil type in both watersheds is muck. The most common soil right around the lake is sand.

Loamy sands tend to be well-drained, with water, air and nutrients moving through them at a rapid rate. Runoff, when it occurs, tends to be slow. Loamy sands have little water-holding capacity and low natural fertility, although they usually have more organic matter present than do sandy soils. Both wind and water erosion are potential hazards with loamy sands, as is drought. There are difficulties with waste disposal and vegetation establishment because of slope and seepage.

Sandy soil tends to be excessively drained, no matter what the slope. Water, air and nutrients move through sandy soils at a rapid rate, so that little runoff occurs unless the soil becomes saturated. Although water erosion can be a problem, wind erosion may be more of a hazard with sandy soils, especially since these soils dry out so quickly. There are also draught hazards with sandy soils. Getting vegetation started in sandy soils is often difficult due to the low available water capacity, as well as low natural fertility and organic material. Onsite waste disposal in sandy soils is also a problem because of slope and seepage; mound systems are usually required.

Land Use in Sherwood Lake Watersheds

Although the ground watershed for Sherwood Lake is fairly small, the surface watershed is quite large. The two most common land uses in the ground watershed are woodlands and residential. The two most common land uses in the surface watershed are woodlands and irrigated agriculture.

Sherwood Lake has a total shoreline 7.8 miles (41,184 feet) Most of the lakeshore is in residential or beach club use. According to a 2004 shore survey, some of the areas near the shore are steeply sloped; some are also soft and/or not well-vegetated. Only 12.84% of the Sherwood Lake shore has native vegetation. 77.54% of the shore has been disturbed and is currently covered by mowed lawn, rock riprap, some kind of seawall, hard structures (piers, etc.), erosion and/or sand.

A 2004 shore survey showed that very little of Sherwood Lake's shore had an "adequate buffer." An "adequate buffer" is a native vegetation strip at least 35 feet landward from the shore. Most of the "inadequate" buffer areas were those with rock or seawall, hard structures, beach, active erosion or mowed lawns. In a few instances, those with insufficient native vegetation at the shoreline to cover 35 feet landward from the water line were also called "inadequate." A vegetated shore is especially important when shores are steep and soft, as are many of Sherwood Lake shores.

Adequate buffers on Sherwood Lake in some places could be easily installed on the inadequate areas by either letting the first 35 feet landward from the water just grow without mowing it, except for a path to the water, or by planting native seedlings sufficient to fill in the first 35 feet or using biologists to protect the shore that are vegetated. Where areas are deeply eroded, shaping, revegetating and protecting the shores will be necessary to prevent further erosion.

Water Testing Results

Between 2004 and 2006, Adams County Land & Water Conservation Department gathered water chemistry and other water quality information on Sherwood Lake. Overall, Sherwood Lake was determined to be a mildly eutrophic lake with fair to poor water quality and poor water clarity.

Measuring the phosphorus in a lake system provides an indication of the nutrient level in a lake. Increased phosphorus in a lake will feed algal blooms and also may cause excess plant growth. The 2004-2006 summer average phosphorus concentration in Sherwood Lake was 37.7 micrograms/liter. This average is over the 30 micrograms/liter level recommended to avoid nuisance algal blooms. This concentration suggests that Sherwood Lake is likely to have nuisance algal blooms from excessive phosphorus.

Water clarity is a critical factor for plants. If plants don't get more than 2% of the surface illumination, they won't survive. Water clarity is measured with a Secchi disk. Average summer Secchi disk clarity in Sherwood Lake in 2004-2006 was 4.36 feet. This is poor water clarity.

Chlorophyll-a concentration provides a measurement of the amount of algae in a lake's water. Algae are natural and essential in lakes, but high algal populations can increase water turbidity and reduce light available for plant growth, as well as result in unpleasing odor and appearance. The 2004-2006 growing season (June-September) average chlorophyll-a concentration in Sherwood Lake was 20.7 micrograms/liter, lower than the state average of 65 micrograms/liter for impoundments.

Sherwood Lake water testing results showed "hard" water with an average of 178 milligrams/liter CaCO₃. Hard water lakes tend to produce more fish and aquatic plants than soft water lakes because they are often located in watersheds with soils that load phosphorus into the lake water.

A lake with a neutral or slightly alkaline pH like Sherwood Lake is a good lake for fish and plant survival. Natural rainfall in Wisconsin averages a pH of 5.6. This means that if the rain falls on a lake without sufficient alkalinity to buffer that acid water coming in by rainfall, the lake's fish cannot reproduce. That is not a problem at Sherwood Lake, since its surface water alkalinity averages 125.6 milliequivalents/liter. The pH levels from the bottom of the lake to the surface hovered between nearly 7 and 8, alkaline enough to buffer acid rain.

Some of the water quality testing at Sherwood Lake showed areas of concern. The presence of a significant amount of chloride over a period of time indicates there may be negative human impacts on the water quality present from septic system failure, the presence of fertilizer and/or waste, deposition of road-salt, and other nutrients. An increased chloride level is thus an indication that too many nutrients are entering the lake, although the level has to be evaluated compared to the natural background data for chloride. The average chloride level found in Sherwood Lake during the 2004-2006 testing period was 11.73 milligrams/liter, elevated substantially above the natural level of 3 milligrams/liter for chloride in this area of Wisconsin. Prior studies also found elevated chloride levels in Sherwood Lake. In fact, substantially elevated chloride levels have been found at Sherwood Lake since records were kept (1985). The source of this ongoing elevation needs to be identified and the elevation reduced.

The sum of water testing results for nitrate, nitrite and ammonium levels of over .3 milligrams/liter in the spring can be used to project the likelihood of an algal bloom in the summer (assuming sufficient phosphorus is also present). Sherwood Lake combination spring levels from 2004 to 2006 averaged .99 milligrams/liter, above the .3 milligrams/liter predictive level for nitrogen-related algal blooms. These elevations suggest that some of the algal blooms on Sherwood Lake may be at least partly nitrogen-related. This issue should be further investigated.

In low-oxygen waters (hypoxic), sulfate can combine with hydrogen and becomes the gas hydrogen sulfate, which smells like rotten eggs and is toxic to most aquatic organisms. Sulfate levels can also affect the metal ions in the lake, especially iron and mercury, by binding them up, thus removing them from the water column. To prevent the formation of hydrogen sulfate, levels of 10 milligrams/liter are best. A health advisory kicks in at 30 milligrams/liter. Sherwood Lake sulfate levels averaged 29.01 milligrams/liter during the testing period, above the level for hydrogen sulfate formation, but still slightly below the health advisory level. However, the overall average for the years in which sulfate testing was done is 30.84, above the health advisory level. This is also an area of concern to be further investigated.

The average calcium level in Sherwood Lake's water during the testing period was 41.36 milligrams/liter. The average Magnesium level was 17.7 milligrams/liter. Both of these are low-level readings. Both sodium and potassium levels in Sherwood Lake are very low: the average sodium level was 3.4 milligrams/liter; the average potassium reading was 2.55 milligrams/liter.

Turbidity reflects water clarity. The term refers to suspended solids in the water column—solids that may include clay, silt, sand, plankton, waste, sewage and other pollutants. Turbid water may mask the presence of bacteria or other pollutants because the water looks murky or muddy. In general, turbidity readings of less than 5 NTU are best. Very turbid waters may not only smell, but also tend to be aesthetically displeasing, thus curtailing recreational uses of the water. Turbidity levels for Sherwood Lake's waters were low in the 1980s, but have risen substantially since, with one reading over the 5 TU mark.

Phosphorus

Like most lakes in Wisconsin, Sherwood Lake is a phosphorus-limited lake: of the pollutants that end up in the lake, the one that most affects the overall quality of the lake water is phosphorus. The amount of phosphorus especially affects the frequency and density of aquatic vegetation and the frequency and density of various kinds of algae, as well as water clarity and other water quality aspects.

The total phosphorus (TP) concentration in a lake is considered a good indicator of a lake's nutrient status, since the TP concentration tends to be more stable than other types of phosphorus concentration. For a man-made lake like Sherwood Lake, a total phosphorus concentration below 30 micrograms/liter tends to result in few nuisance algal blooms. Sherwood Lake's growing season (June-September) surface average total phosphorus level of 37.7 micrograms/liter is over that limit, suggesting that phosphorus-related nuisance algal blooms may occur.

Land use plays a major role in phosphorus loading. The land uses around Sherwood Lake that contribute the most phosphorus are irrigated and non-irrigated agriculture. Some phosphorus deposition cannot be controlled by humans. However, some phosphorus (and other nutrient) input can be decreased or increased by changes in human land use patterns. Practices such as shoreland buffer restoration along waterways; infiltrating stormwater runoff from roof tops, driveways and other impervious surfaces; using no phosphorus lawn fertilizers; and reducing phosphorus input to and properly managing septic systems will minimize phosphorus inputs into the lake. Such practices need to be implemented in all of the Tri-Lakes Watersheds in order for a significant impact on phosphorus reduction to occur.

Reducing the amount of input from the surface and ground watersheds results in less nutrient loading into the lake itself. Under the modeling predictions, reducing phosphorus inputs from human-based activities even 10% would improve Sherwood Lake water quality by 1 to 8.6 micrograms/liter. A 25% reduction would save 2.5 to 21.5 micrograms/liter and could reduce the overall epilimnetic growing season total phosphorus to around the 30 micrograms/liter level to avoid nuisance algal blooms. These predictions make it clear that reducing current phosphorus inputs to the lake are essential to improve, maintain and protect Sherwood Lake's health for future generations.

Aquatic Plant Community

The aquatic plant community is characterized by below average quality for Wisconsin lakes, poor species diversity and impacted by high levels of disturbance. Sherwood Lake is within the 25% of lakes in the state most tolerant of disturbance and furthest from an undisturbed condition. Disturbances include invasions of exotic species, boat traffic, shoreline development, harvesting and past herbicide treatments.

Of the 29 species found in Sherwood Lake, 25 were native and 4 were exotic invasives. In the native plant category, 14 were emergent, 1 was a free-floating plant, and 10 were submergent species. Four exotic invasives, *Myriophyllum spicatum* (Eurasian Watermilfoil), *Nasturtium microphyllum* (watercress), *Phalaris arundinacea* (Reed Canarygrass), and *Potamogeton crispus* (Curly-Leaf Pondweed) were found. Filamentous algae were found at 22.76% of the sample sites in 2006 and at 38.33% of the sites in 2000.

Potamogeton pectinatus, an aquatic plant favored by drawdowns, was the most frequently-occurring plant in Sherwood Lake in 2006. In 2000, the most frequent species was *Chara* spp. No species but *Potamogeton pectinatus* reached a frequency of 50% or greater in the lake overall in 2006, although *Chara* spp and *Potamogeton crispus* were not far under 50%, with occurrence frequencies of 45.53% and 42.28%

respectively. In 2000, no aquatic species reached an overall occurrence frequency of over 50%.

Potamogeton pectinatus was also the densest plant in 2006 in Sherwood Lake. In the lake overall, none of the aquatic plant species had a mean density of over 2.0, meaning none occurred at more than average, in 2006. In 2006, the only species occurring at more than average density in any of the depth zones was *Potamogeton pectinatus* in the second (1.5 feet-5 feet) and third (5 feet-10 feet) depth zones. Densest in Depth Zone 1 (0 to 1.5 feet) was *Chara spp*; densest in the other three zones was *Potamogeton pectinatus*. No species occurred at more than average density in the lake overall in 2000, either. The only depth zone with more than average density of growth was Depth Zone 3, where *Chara spp* grew at more than average density.

Relative frequency and relative density are combined into a dominance value that demonstrates how dominant a species is within its aquatic plant community. Based on dominance value, *Chara spp* was the dominant aquatic “plant” species in Sherwood Lake in 2000. Sub-dominant was *Elodea canadensis*. However, in 2006, *Potamogeton pectinatus* dominated the aquatic plant community, with *Potamogeton crispus* and *Chara spp* next most dominant.

Looking at the results from the 2000 survey and those from 2006 shows some changes in the aquatic plant community. There were more species found in 2006, and the structure of the aquatic plant community has changed with more emergent species present, but only one free-floating plant. No floating-leaf plants, which provide habitat and cover for fish and invertebrates, were found in either year. Further, when calculating the coefficient of similarity between the 2000 and 2006 surveys, they score as statistically dissimilar both in terms of frequency of occurrence and relative frequency. Based on frequency of occurrence, the aquatic plant communities of the two years are just over 45% similar. Using relative frequency, the score is only 53% similar. Similarity percentages of 75% or more are considered statistically similar. Obviously, the figures for Sherwood Lake are far below that figure.

It is worth noting that the report on the 2000 aquatic plant surveys mentioned the low level of emergent plants in Sherwood Lake. The 2006 survey shows that emergent plants were still scarce in Sherwood Lake than they were in 2000, but there were more increased coverage from emergent plants in 2006.

Sherwood Lake has five known invasive aquatic plant species: Curly-Leaf Pondweed (submergent); Eurasian Watermilfoil (submergent); Purple Loosestrife (emergent); Reed Canarygrass (emergent) and Watercress (floating-leaf). The lake gets a significant amount of transient boat traffic due to its location (right off a main highway) and large public boat ramp. The Tri-Lakes Management District has a lake

management plan that includes management of aquatic invasives. The lake has been using targeted harvesting for Eurasian Watermilfoil, emphasizing the harvesting of that plant in May and September, while harvesting the summer months for navigation, rather than control of Eurasian Watermilfoil. In 2007, some lake citizens were trained to monitor the aquatic invasives and participate in the Clean Boats, Clean Waters boater education program.

Fish/Wildlife/Endangered Resources

WDNR stocking and fishery inventories go back to 1968, when the lake was stocked after a chemical eradication of fish in 1967 to get rid of the rough fish population. Stocking in 1968 consisted of bluegills, largemouth bass, northern pike and walleye. A follow-up inventory in 1969 found that bluegills and pumpkinseeds were abundant; largemouth bass, northern pike, walleye and yellow perch were common; and shiners and white suckers were scarce. The most recent survey, done in 2002, found that bluegills and largemouth bass were abundant; black crappie, walleye and yellow perch were common; and northern pike was scarce. Between 1970 and 2000, thirteen other fish inventories were performed by the WDNR. In addition to those fish already mentioned, various fish surveys through the years also found brown bullheads, black bullheads, yellow bullheads, yellow suckers, golden shiners, and emerald shiners.

In 1999, the local WDNR fishery biologist reported that a recent survey of Sherwood Lake showed that the largemouth bass and northern pike populations were good and that those fish were healthy. However, although the panfish numbers were sufficient, he found them to be small and thin. He expressed concern about loss of invertebrate habitat that the fish fed on, about the chemicals killing zooplankton, and about the effects of the winter drawdown on the aquatic plant community.

There are several endangered resources in the Sherwood Lake surface watershed. Natural communities reported here include northern sedge meadow, northern wet forest, pine barrens and shrub-carr. Endangered plants known in the area include *Polygala cruciata* (crossleaf milkwort), *Juncus marginata* (grassleaf rush), and *Bartonia virginica* (yellow screwstem).

Conclusion

Sherwood Lake is currently impoundment impacted substantially by its position in the large surface watersheds of the Tri-Lakes, as well as significant disturbances. It is approaching the threshold of passing from an aquatic plant-dominated system to a turbid algae-dominated system. The Tri-Lakes Management District will need to regularly review and update its lake management plan in order to address its management issues in a logical, cohesive manner.

RECOMMENDATIONS

Lake Management Plan

- When the Tri-Lakes Management District revises the lake management plan, it needs to make sure the plan includes at least the following aspects concerning the management of the lake: integrated aquatic species management; control/management of invasive species; wildlife and fishery management; nutrient budgeting; shoreland protection; water quality protection.
- The Sherwood Property Owner's Association should participate in the revision process and implementation of the lake management plan.

Watershed Recommendations

- Since computer modeling results suggest that input of nutrients, especially phosphorus, are a factor that needs to be explored for Sherwood Lake, it is recommended that both the surface and ground watersheds be inventoried, documenting any of the following: runoff from any livestock operations that may be entering the surface water; soil erosion sites; agricultural producers not complying with nutrient management plans and/or irrigation water management plans.
- If such sites are documented, a statement outlining the Sherwood Lake Association and Tri-Lake Management District's encouragement to Adams County Land & Water Conservation Department and landowners to design and implement practices to address the sites.

Water Quality Recommendations

- All lake residents should practice best management on their lake properties, including keeping septic systems maintained in proper condition and pumped every three years, eliminating the use of lawn fertilizers, cleaning up pet wastes and not composting near the water.
- Reducing the amount of impervious surface around the lake and management of stormwater runoff will also help maintain water quality.
- Residents should become involved in the Citizen Lake Water Monitoring Program. This program includes water quality monitoring, invasive species monitoring, and Clean Boats, Clean Waters.
- Broad-scale restoration of native vegetation at the shore is needed to help improve water quality. Studies show that the frequency and density of the most sensitive plant species is less at disturbed shores than at those with native vegetation. These sensitive plants are indicators of changing water quality.

- Further investigation of the sources of the elevated chloride, nitrogen and sulfate needs to be made to identify such sources and develop a plan to reduce those elevated levels.

Aquatic Plant Recommendations

- All lake users should protect the aquatic plant community in Sherwood Lake by assisting in revising implementing an integrated aquatic plant management plan that uses multiple methods of control.
- The Tri-Lakes Management District should maintain exotic species signs at the boat landings and contact DNR if the signs are missing or damaged.
- The Tri-Lakes Management District should continue monitoring and control of Eurasian Watermilfoil and Curly-Leaf Pondweed, maintaining the most effective methods and modifying if necessary. The Sherwood Property Owners Association should assist in these efforts. Residents may need to hand-pull scattered plants.
- Lake residents should get involved in the county-sponsored Citizen Aquatic Invasive Species Monitoring Program. This will allow not only noting changes in the Eurasian Watermilfoil pattern and Curly-Leaf Pondweed, but also for other invasives, including the zebra mussels already known to be present in Arrowhead Lake. Noting the presence and density of invasives early is the best way to take preventive action to keep them from becoming a bigger problem.
- Emergent vegetation, which is very sparse in Sherwood Lake, should not be removed. In fact, removal of aquatic plants and shore plants should be kept to the maximum 30' wide viewing/access corridor and navigation purposes. Leave as much vegetation as possible to protect water quality and habitat.
- Natural shoreline should be restored and eroded areas repaired. Disturbed shoreline covers too much of the shore and mowed lawn alone covers nearly half of the shore.
 - a) Unmowed native vegetation reduces shoreline erosion and run-off into the lake and filters the run-off that does enter the lake thus reducing nutrient inputs.
 - b) Shoreline restoration could be as simple as leaving a band of natural vegetation around the shore by discontinuing mowing.
 - c) Restoration could be as ambitious as extensive plantings of attractive native wetland species in the water and native grasses, flowers, shrubs and trees on the near shore area.

LAKE CLASSIFICATION REPORT FOR SHERWOOD LAKE, ADAMS COUNTY

INTRODUCTION

In 2003, The Adams County Land & Water Conservation Department (Adams County LWCD) determined that a significant amount of natural resource data needed to be collected on the lakes with public access in order to provide it and the public with information necessary to manage the lakes in a manner that would preserve or improve water quality and keep it appropriate for public use. In some instances, there was significant historical data about a particular lake; in that instance, the study activities concentrated on combining and updating information. In other instances, there was no information on a lake, so study activities concentrating on gathering data about that lake. Further, it was discovered that information was scattered among various citizens, so often what information was actually available regarding a particular lake was unknown. To assist in updating some information and gathering baseline information, plus centralize the data collected, so the public may access it. The Adams County LWCD received a series of grants from the Wisconsin Department of Natural Resources (WDNR) from the Lake Classification Grant Program.

Objectives of the study were:

- collect physical data on the named lakes to assist in assessing the health of Adams County lake ecosystems and in classifying the water quality of the lakes.
- collect chemical and biological data on the named lakes to assist in assessing the health of Adams County lake ecosystems and in classifying the water quality of the lakes.
- develop a library of lake information that is centrally located and accessible to the public and to City, County, State and Federal agencies.
- make specific recommendations for actions and strategies for the protection, preservation and management of the lakes and their watersheds.
- create a baseline for future lake water quality monitoring.
- Provide technical information for the development of comprehensive lake management plans for each lake
- provide a basis for the water quality component of the Adams County Land and Water Resource Management Plan. Components of the plan will be incorporated into Adams County's "Smart Growth Plan".
- develop and implement educational programs and materials to inform and educate lake area property owners and lake users in Adams County.

METHODS OF DATA COLLECTION

To collect the physical data, the following methods were used:

- delineation & mapping of ground & surface watersheds using topographic maps, ground truthing and computer modeling;
- identification of flow patterns for both the surface & ground watersheds using known flow maps and topographic maps;
- inventory & mapping of current land use with orthographic photos and collected county information;
- inventory & mapping of shoreline erosion and buffers using county parcel maps and visual observation;
- inventory & mapping for historical and cultural sites using information from the local historical society and the Wisconsin Historical Society;
- identification & mapping of critical habitat areas with WDNR and Adams County LWCD staff;
- identification & mapping of endangered or threatened natural resources (including natural communities, plant & animal species) using information from the Natural Heritage Inventory of Wisconsin;
- identification & mapping of wetland areas using WDNR and Natural Resource Conservation Service wetland maps;
- preparation of soil maps for each of the lake watersheds using soil survey data from the Natural Resource Conservation Service.

To collect water quality information, different methods were used:

- for three years, lakes were sampled during late winter, at spring and fall turnover, and several times during the summer for various parameters of water quality, including dissolved oxygen, relevant to fish survival and total phosphorus, related to aquatic plant and algae growth;
- random samples from wells in each lake watershed were taken in two years and tested for several factors;
- aquatic plant surveys were done on all 20 lakes and reports prepared, including identification of exotics, identifying existing aquatic plant community, evaluation of community measures, mapping of plant distribution, and recommendations;
- all lakes were evaluated for critical habitat areas, with reports and recommendations being made to the respective lakes and the WDNR;
- lake water quality modeling was done using data collected, as well as historical data where it was available.

WATER QUALITY COMPUTER MODELING

Wisconsin developed a computer modeling program called WiLMS (Wisconsin Lake Modeling Suite) to assist in determining the amount of phosphorus being loaded annually into a lake, as well as the probable source of that phosphorus. This suite has many models, including Lake Total Phosphorus Prediction, Lake Eutrophic Analysis Procedure, Expanded Trophic Response, Summary Trophic Response, Internal Load Estimator, Prediction & Uncertainty Analysis, and Water & Nutrient Outflow. The models that various types of data inputs: known water chemistry; surface area of lake; mean depth of lake; volume of lake; land use types & acreage. This information is then used in the various models to determine the hydrologic budget, estimated residence time, flushing rate, and other parameters.


Using the data collected over the course of the studies, various models were run under the WiLMS Suite. These water quality models are computer-based mathematical models that simulate lake water quality and watershed runoff conditions. They are meant to be a tool to assist in predicting changes in water quality when watershed management activities are simulated. For example, a model might estimate how much water quality improvement would occur if watershed sources of phosphorus inputs were reduced. However, it should be understood that these models predict only a relative response, not an exact response. Modeling results will be incorporated into topic discussions as appropriate.

DISSEMINATION OF PROJECT DELIVERABLES

The results of this study will be distributed to various agencies, organizations and the public as previously described. Based on the classification information, the Adams County Land and Water Conservation Department will identify assistance requests and determine the appropriate future activities, based on the classification determinations. To provide the requested assistance, Adams County Land and Water Conservation Department will incorporate the lake management plans goals, priorities and action items into its Annual Plan of Operations. Goals, priorities and action items may include educational programs, formation of lake districts, further development of lake management plans and implementation of lake management plans.

ADAMS COUNTY INFORMATION

Adams County lies in south central Wisconsin, shaped roughly like the outline of Illinois. Adams County is a small rural county with a full-time population of about 20,000. Between 1980 and 2000, Adams County's population grew by more than 20%, with most of the population increase being located upon the lakes and streams. The population increase has resulted in a greater need for facilitation, technical assistance and education, including information on the lakes and streams.




**Figure 1:
Adams
County
Location in
Wisconsin**

SHERWOOD LAKE BACKGROUND INFORMATION

Sherwood Lake is located in the Town of Rome, Adams County, WI, in the south central part of Wisconsin. The impoundment of 14-Mile Creek is slightly over 243 surface acres in size. Maximum depth is 24', with an average depth of 8'. Both Upper and Lower Camelot Lakes flow through dams into Sherwood Lake. Sherwood Lake flows through a dam into Arrowhead Lake. All the Tri-Lakes dams are owned and operated by Adams County. There is a public boat launch on Sherwood Lake on the southwest edge of the lake owned by the Parks Department of Adams County. Heavy residential development around the lake is found along most of the lakeshore. Sherwood Lake is managed by the Tri-Lakes Management District. There is also an active Sherwood Property Owners Association.

Figure 2: Sherwood Lake location




RE:2/07

The Central Sand Plains, which contain Sherwood Lake, are found in the Driftless Area of Wisconsin. The area is characterized by varying elevations, with numerous, usually flat-topped ridges & hills sometimes called “mounds.” Deposits made by streams from the melting ice sheet cover large areas and usually consist of sand, clay and gravel.

Archeological Sites

Figure 3: Sherwood Lake Archeological Sites



RE:4/05: revised 7/06

There are many Native American archeological sites in Adams County, with several being located right around in the Tri-Lakes watersheds. These mounds can be conical, linear or effigy (animal shapes) shapes. In order to preserve Native American heritage, federal and state laws on Native American burials require permission of the federal government and input from the local tribes before further disturbance.

Bedrock and Historical Vegetation

Bedrock around Sherwood Lake is mostly sandstone, both weak and resistant, formed in the Cambrian Period of Geology (542 to 488 millions years ago). Bedrock may be 200 or more feet below the sand/clay/gravel deposits left by melting ice cover.

Original upland vegetation of the area included extensive wetlands of many types (including open bogs, shrub swamps & sedge meadows), as well as prairies, oak forests, savannahs and barrens. Mesic white pine & hemlock forests were found in the northwest portion of the region. Most of the historic wetlands were drained in the 1900s and used for cropping. The current forested areas are mostly oak-dominated, followed by aspen and pines. There are also small portions of maple-basswood forest and lowland hardwoods.

Soils in the Sherwood Lake Watersheds

The primary soil type in both the surface and ground watersheds is loamy sand. The second most common soil type in both watersheds is muck. The most common soil right around the lake is sand.


Loamy sands tend to be well-drained, with water, air and nutrients moving through them at a rapid rate. Runoff, when it occurs, tends to be slow. Loamy sands have little water-holding capacity and low natural fertility, although they usually have more organic matter present than do sandy soils. Both wind and water erosion are potential hazards with loamy sands, as is drought. There are difficulties with waste disposal and vegetation establishment because of slope and seepage.

Sandy soil tends to be excessively drained, no matter what the slope. Water, air and nutrients move through sandy soils at a rapid rate, so that little runoff occurs unless the soil becomes saturated. Although water erosion can be a problem, wind erosion may be more of a hazard with sandy soils, especially since these soils dry out so quickly. There are also draught hazards with sandy soils. Getting vegetation started in sandy soils is often difficult due to the low available water capacity, as well as low natural fertility and organic material. Onsite waste disposal in sandy soils is also a problem because of slope and seepage; mound systems are usually required.

The soil and soil slopes around lakes and streams are very important to water quality. They affect amount of infiltration of surface precipitation into the ground and the amount of contaminants that may reach the groundwater, as well as the amount of

surface stormwater runoff. In addition, these two factors affect the amount and content of pollutants and particles (including soil) that may wash into a water body, affecting its water quality, its aquatic plant community and its fishery. Further, soil types and soil slopes help determine the appropriate private sewage system and other engineering practices for a particular site, since they affect absorption, filtration and infiltration of contamination from engineering practices.

Figure 4: Sherwood Lake Soils



RE:2/05

PRIOR STUDIES OF THE TRI-LAKES AREA

The Tri-Lakes Area, including Sherwood Lake, has been the subject of several different studies. In 1978, a “water management study” report, Fourteenmile Creek Watershed, was published by the University of Wisconsin-Madison. This summarized the results of a study done during the summer of 1978. The report noted that the first lake constructed in the system was Sherwood Lake, on which construction started in 1967. By 1978, all of the Tri-Lakes had been constructed. The upper watershed was used mostly for agriculture and commercial forest. 70% of the agricultural fields were irrigated. Because of the concerns expressed by the landowners in the lower watershed (around the lakes), the report concentrated on studying that area.

A survey of lakeshore owners indicated that water weeds were seen as the most severe problem, followed by erosion, algal blooms and oil slicks from boat motors. There were also complaints about water “muddiness.” Over one-half of the respondents thought that contributors to these problems included leaking septic systems, oil from boat motors, silt & sand from shoreline erosion, construction erosion and fertilizers from lawns and fields. Researchers saw lakes Camelot and Sherwood as more vulnerable to eutrophy because of their shallowness, relative clarity and abundant nutrients to encourage dense aquatic plant and algal growth. Sherwood Lake was determined to be fed by water over Camelot Lake dams and groundwater. Because Sherwood Lake waters mixed frequently, nutrients from the bottom of the lake was made available to algae and plants in the surface waters. Sherwood Lake also had a high flushing rate, with its total volume of water being replaced about 15 times per year. High flushing rates often bring greater loads of nutrients. The researchers noted that substantial shoreline erosion was occurring on Sherwood Lake. They indicated that the main causes of erosion were waves stirred up by winds or boat wakes. Further study revealed that wind was responsible for three times as many waves as boat wakes, but that boat wakes created many more large waves.

Recommendations from this report included: (1) installation of shoreline protection practices to reduce erosion; (2) lakeside vegetation planting to reduce soil loss and help control erosion; (3) amendment of the Town of Rome boating ordinance to reduce the incidence of high energy, boat-generated waves by specifying speeds without certain distances from shores; (4) better enforcement of the current town boating ordinance to control boat speeds and water skiing issues; (5) continued stocking of game fish like pike & walleye, as well as fish habitat improvement within the lakes; (6) establishing a lake management district that included all the lakes.

In 1993, another report called Tri-Lakes, Adams County, Wisconsin: Lakes and Watershed Characterization was published. This study was funded by the WDNR and the Tri-Lakes Management District and performed by Blue Water Science of St Paul, Minnesota. Parameters researched during this study include collection of flow & total phosphorus data for the creek and ditches feeding into the Tri-Lakes; collecting lake water chemistry sample for one summer (1991); and estimating phosphorus load from various land uses and contribution sources. Runoff was estimated at 6.3 inches per year. Secchi disk readings for Sherwood Lake ranged from 10.5 feet (June 1991) to 4 feet (August 1991). Hypoxia in the water column increased as the summer went on. In June, hypoxia was present only at 25 feet or more; by August, hypoxia started at about 8 feet in depth. Total phosphorus for summer 1991 ranged from 20 to 32 micrograms/liter. Chlorophyll-a readings ranged from 6 to 21 micrograms/liter. Although total phosphorus levels were generally low, nitrate levels were always very high. An overall trophic state index of 52 was calculated for Sherwood Lake. These researchers calculated an overall total phosphorus load of nearly 3740 pounds per year to Sherwood Lake.

The report made several recommendations; (1) to continue efforts to reduce wind erosion through practices such as tree planting, snow fence planting, residue management, etc.; (2) to continue monitoring incoming stream water and lake water, examining at least total phosphorus, nitrate-nitrite, total Kjeldahl nitrogen and total suspended solids; (3) conduct bioassays on the 14-Mile Creek bedloads, since it was estimated that their phosphorus level was considerably higher than that of the creek surface water; (4) to continue a lake and watershed information & education program using activities such as newsletters, lake fair picnics, underwater video, demonstration projects, appointment of lake captains responsible for information distribution on their lakes; (5) to conduct plan surveys & monitor the amount of plant removal by monthly analyzing plants mechanically harvested for total phosphorus on each lake; (6) to install demonstration projects for aquatic plant control nearshore by methods other than herbicides; (7) to landscape lake shores for wildlife, shore protection & erosion control; and (8) to perform an on-site waste system evaluation and conductivity study, taking samples that were analyzed for fecal coliform & fecal streptococcus bacteria.

In April, 1999, Mid-State Associates Professional Services published the results of a septic study it had done. 45% if the properties around Sherwood Lake were developed at that time. Houses tended to be within 100 feet of the waterfront, with lakeshores narrow and houses close together. MSA inspected 20% of the septic sites. It found that 18% of the septic systems around Sherwood Lake were failing. While the overall average age of Sherwood Lake's septic systems was 18 years, the average age of the failing systems was 27 years. The report indicated that up to 31% of the septic systems over 20 years old would up expected to fail, with effects most obvious in localized

sections of the lake. It also noted that the sandy soils in the drainfields had a low capacity to retain phosphorus. According to this report, previous studies in 1979 and 1993 had overestimated the sandy soils' ability to retain phosphorus. The report also indicated that septic systems phosphorus input was likely to grow as development increased and the sandy soils reached their phosphorus saturation point. The report recommended four alternatives to be considered in place on continuing to use many individual private septic systems: (1) centrally collect wastewater and discharge it outside the Tri-Lakes watersheds; or (2) centrally collect wastewater and pump it into an existing municipal sewage system; or (3) use cluster type wastewater collection with nutrient removal and discharge of treated water; or (4) use nutrient removal techniques in the individual on-site waste systems.

The results of a study of the algae in Sherwood Lake were outlined in a report titled Phytoplankton Community Composition and Distribution in the Tri-Lakes Area, written by Dr. Robert Bell, UWSP-Biology Department, published in 2000. He found that Sherwood Lake had the most taxonomically diverse algae of all the Tri-Lakes. However, the taxa were generally unremarkable and were seen as typical of a mesotrophic or slightly eutrophic lake. Dr. Bell felt that at that time, heavy aquatic plant growth was more of a problem than seasonal algal blooms, but that he expected the levels of cyanobacteria (blue-green algae) to increase as the algal community shifted from one roughly equally spot of cyanobacteria, ochrophytes (pigmented algae) and green algae to one of predominately cyanobacteria. He made the following recommendations: (1) reduce the upstream input of pesticides & growth-promoting nutrients by using sediment traps or lagoons; (2) remove in-lake nutrients via sediment and/or plant biomass removal; (3) reduce residential nutrient input by improved septic/sanitation systems and shoreline vegetation filter strips.

In December 2001, a report titled Assessment of Shallow Groundwater Flow & Chemistry & Interstitial Water Sediment, Aquatic Macrophyte and Chemistry for the Tri-Lakes, Adams County, WI, was published. It was written by B. Shaw, C. Sparacio, J. Stelzer and N. Turyk of UWSP. Objectives of this study were: (1) to compare groundwater flow patterns during full & drawn-down conditions; (2) to examine water quality after heavy summer use; (3) to monitor groundwater entering back bays in Camelot & Sherwood Lakes for local impacts such as septic and/or lawns; (4) to determine nutrients & biomass of aquatic macrophytes as they relate to nutrients in interstitial water & lake sediments; and (5) to determine quantity of phosphorus and nitrogen held by plant tissues to estimate harvest removal. This study found that during non-drawdown conditions, 47% of Sherwood Lake sites showed upwelling, i.e., showed that groundwater was entering the lake. The main downwelling was found near the dam, which is typical. During a drawdown, even more of the sample sites were upwelling, raising the potential for effect on water quality by the water being drawn into the lake. 15.2% of the upwelling sites showed

evidence of septic contamination. 27.3% of these sites showed evidence of negative impacts from land use. 6.1% showed evidence of contamination from road-salt. When the study looked at aquatic plants (macrophytes) and the harvesting done by the Tri-Lakes Management District, it noted that 44% of the harvested tonnage came from Sherwood, with July being the peak month. The study concluded that the fall drawdown & spring refill of Sherwood Lake and the Camelot Lakes resulted in nutrients being released from anaerobic sediments and negatively impacted lake water quality. The high reactive phosphorus and ammonium levels suggested that nutrient transport to the lake was significant, especially from the fall drawdown, which increased groundwater discharge to the lake. The report stated that there was nutrient availability from several sources, including (1) the anaerobic release of ammonium and phosphorus from high organic matter sediments; (2) the nutrient flux from groundwater inflow; and (3) the cycling of nutrients from lake to sediment to groundwater and back again.

A survey was done of Tri-Lakes property owners during 2001. Although 72% of the respondents felt that the lakes' water quality was "good" or "fair", 74% felt the water quality had declined since they started coming to the lake. The top three causes attributed by respondents were input from cranberry growers, fertilizer use and heavy recreational use. The presence of algal scum and reduced water clarity were cited as the top two water quality problems, with aquatic plant growth scoring near the bottom. Nearly 66% of the respondents fertilized their lawn, with only 25% using non-phosphorus fertilizer. 74% mowed over 25% of their lawn at their lakefront. 83.8% of the houses had 2 or 3 bedrooms. Main activities on the lake included swimming, boating and fishing. The three most common boat types were pontoon, fishing and skiing, with 56.7% of the respondents having a boat motor 50 horsepower or larger.

A report titled Limnological Investigations of Camelot, Sherwood and Arrowhead Lakes, WI, was written by W.F. James, J. Barko & H. Eaken of the U.S. Army Corps of Engineers in 2002. Field work for this report included evaluating total sediment, total nitrogen and total phosphorus loads in the lakes. This was done by sediment collecting & testing, as well as water quality monitoring and computer modeling. The report noted that total phosphorus and chlorophyll-a levels increased as one went west in the lake chain, with the Camelot Lakes having the lowest levels and Arrowhead Lake having the highest level, with Sherwood Lake between the two. Summer anoxia in the lower levels of the lake was noted in July and August, especially near the Sherwood Dam. Secchi disk readings were lowest in August and September. The average summer Secchi disk reading was 7.2 feet. Average total phosphorus was 15 micrograms/liter and average chlorophyll-a was 14.8 micrograms/liter. The report indicated that Sherwood had an intermediate algal bloom frequency. Report authors determined Sherwood Lake to be susceptible to declining water quality as phosphorus loading increased.

CURRENT LAND USE

Although the ground watershed for Sherwood Lake is fairly small, the surface watershed is quite large. The two most common land uses in the ground watershed are woodlands and residential. The two most common land uses in the surface watershed are woodlands and irrigated agriculture. (See Figures 5 through 8).

Figure 5: Sherwood Lake Watersheds Land Use in Acres and Percent of Total

	Ground		Surface		Total	
Sherwood Lake	Acres	% of Total	Acres	% of Total	Acres	% of Total
Agriculture--Non Irrigated	0	0.00%	8598.14	16.93%	8598.14	16.11%
Agriculture--Irrigated	0	0.00%	14,712.35	28.97%	14712.35	27.57%
Grassland/Pasture	173.89	6.73%	3983.82	7.85%	4157.71	7.79%
Residential	819.83	31.73%	2,627.44	5.17%	3447.27	6.46%
Water	347.78	13.46%	603.9	1.19%	951.68	1.78%
Woodland	1242.2	48.08%	20,252.97	39.89%	21495.17	40.28%
total	2583.7	100.00%	50778.62	100.00%	53362.32	100.00%


Prior information on the watersheds shows how land use has changed over the years. After a substantial increase in agricultural land use between 1978 and 1986, agricultural changes appear to have leveled off and are no longer increasing in acreage. Residential use in the watersheds has decreased overall, although residential use directly around the Tri-Lakes has increased. Woodlands have increased slightly.

Figure 6: Prior Land Use Data for Sherwood Lake Watersheds

Land Use Type	1978	1986	1998	2004
Agricultural (both types)	31.1%	48.6%	48.9%	43.7%
Pasture/Grassland	17.1%	11.3%	11.1%	7.8%
Residential	12.4%	7.8%	9.3%	6.5%
Woodland	37.8%	30.7%	29.1%	40.3%
Water	1.6%	1.6%	1.6%	1.8%

Studies have shown that land use around a lake has a great impact on the water quality of that lake, especially in the amount and content of surface runoff. (James, T., 1992, I-10; Kibler, D.F., ed. 1982. 271) For example, while natural woodland may (on the average) absorb 3.5” out of a 4” rainfall, leaving only .5” as runoff, a residential area with quarter-acre lots may absorb only 2.3” of the 4”, leaving 1.7” to run off the land into the lake—the same amount as may be expected to run off from a corn or soybean field. 1.7” of runoff translates into 46,200 gallons per acre ending up in the lake! Percentage of impervious surface, the soil type, vegetation present and slope of the site can all affect runoff volume. (Frankenberger, J, ID-230). The changes in the Sherwood watershed land use are therefore likely to significantly increase the runoff in volume and content unless protection steps are taken.

Figure 7a: Land use in Sherwood Lake Surface Watershed



Land Use


- NON-IRRIGATED AGRICULTURE
- IRRIGATED AGRICULTURE
- GOVERNMENTAL/COMMERCIAL
- GRASSLAND/PASTURE
- OUTDOOR RECREATION
- RESIDENTIAL
- WATER
- WETLANDS
- WOODLANDS

Surface Watershed Boundary



Land Use--Sherwood Lake Surface Watershed

RE:2004



Land Use in Sherwood Lake Ground Watershed

Land Use



- NON-IRRIGATED AGRICULTURE
- IRRIGATED AGRICULTURE
- GOVERNMENTAL/COMMERCIAL
- GRASSLAND/PASTURE
- OUTDOOR RECREATION
- RESIDENTIAL
- WATER
- WETLAND
- WOODLANDS

Ground Watershed Boundary

RE:2004

Figure 7b: Land Use in Sherwood Lake Ground Watershed

When water runs over a surface, it picks up whatever loose pollutants—sediment, chemicals, metals, exhaust gas, etc—are present on that surface and takes those items with it into the lake. Increased development around a lake tends to increase the amount of pollutants being carried into the lake, thus negatively affecting water quality. Residential development areas with lots of one-quarter acre or less may deliver as much as 2.5 pounds of phosphorus per year to the lake for each acre of development.



There are two specific kinds of land use—wetlands and shorelands--that are so important to water quality that they will be separately discussed.

WETLANDS

A number of wetlands are located in the Sherwood Lake surface and ground watersheds. In the past, wetlands were seen as “wasted land” that only encouraged disease-transmitting insects. Many wetlands were drained and filled in for cropping, pasturing, or even residential development. In the last few decades, however, the importance of wetlands has become evident, even as wetlands continue to decline in acreage.

Wetlands play an important role in maintaining water quality by trapping many pollutants in runoff and flood waters, thus often helping keep clean the water they connect to. They serve as buffers to catch and control what would otherwise be uncontrolled water and pollutants. Wetlands also play an essential role in the aquatic food chain (thus affecting fishery and water recreation), as well as serving as spaces for wildlife habitat, wildlife reproduction and nesting, and wildlife food.


The large areas of wetlands in the Sherwood Lake watersheds serve as filters and traps that help keep Sherwood Lake as clean as it is. It is essential to preserve these wetlands for the health of all the Tri-Lakes.



Figure 9:
Example of
lake end
wetland

SHORELANDS

Sherwood Lake has a total shoreline 7.8 miles (41,184 feet) Most of the lakeshore is in residential or beach club use. According to a 2004 shore survey, some of the areas near the shore are steeply sloped; some are also soft and/or not well-vegetated. Only 12.84% of the Sherwood Lake shore has native vegetation. 77.54% of the shore has been disturbed and is currently covered by mowed lawn, rock riprap, some kind of seawall, hard structures (piers, etc.), erosion and/or sand.




Some of the Sherwood Lake shores were heavily eroded, as shown in Figure 11.



**Figure 11:
Example of Erosion
on Sherwood Lake**

Sherwood Lake Shoreline



- Active Erosion
- Beach or Sand
- Rock or Seawall
- Vegetated Shore



Figure 12: Shoreland Map of Sherwood Lake (2004)

The Adams County Shoreline Ordinance defines 1000' landward from the ordinary high water mark as "shoreland". Under the ordinance, the first 35 feet landward from the water is a "buffer." Shoreland buffers are an important part of lake protection and restoration. These buffers are simply a wide border of native plants, grasses, shrubs and trees that filter and trap soil & similar sediments, fertilizer, grass clippings, stormwater runoff and other potential pollutants, keeping them out of the lake. A 1990 study of Wisconsin shorelines revealed that a buffer of native vegetation traps 5 to 18 times more volume of potential pollutants than does a developed, traditional lawn or hard-armored shore.

A 2004 shore survey showed that very little of Sherwood Lake's shore had an "adequate buffer." An "adequate buffer" is a native vegetation strip at least 35 feet landward from the shore. Most of the "inadequate" buffer areas were those with rock or seawall, hard structures, beach, active erosion or mowed laws. In a few instances, those with insufficient native vegetation at the shoreline to cover 35 feet landward from the water line were also called "inadequate."

Vegetated shoreland buffers help stabilize shoreline banks, thus reducing bank erosion. The plant roots give structure to the bank and also increase water infiltration and decrease runoff. A vegetated shore is especially important when shores are steep and soft, as are many of Sherwood Lake shores. Figure 13 maps the adequate and inadequate buffers on Sherwood Lake.

Sherwood Lake--Buffers



 Adequate Buffer  Inadequate Buffer



RE:2004

Figure 13: Sherwood Lake Buffer Map (2004)

Lakeside buffers also serve as important habitat. Lake edges usually contain aquatic and wetland plants, grading into drier groundcover, then shrubs and trees as one moves inland towards drier land. Buffers provide habitat for many species of water-dependent wildlife, including furbearers, reptiles, birds and insects. Many wildlife species, including birds, small mammals, fish & turtles breed, nest, forage and/or perch in shore buffer areas. Further, 80% of the endangered and threatened species listed spend part of their life in this near-lake buffer area. (Wagner et al, 2006)

When the natural shoreline is replaced by traditional mowed turf-grass lawns, rock, wooden walls or similar installments, bird and animal life, land-based insects, and aquatic insects that hatch or winter on natural shore are negatively impacted. For example, on many Adams County lakes, the non-native aquatic plant, Eurasian Watermilfoil has invaded. There is a weevil native to Wisconsin that weakens Eurasian Watermilfoil by burrowing into and developing within its stems, but that weevil depends on a native-plant shore to overwinter. If the shore is instead covered by rock, seawall or traditional lawn, these weevils will be unavailable for the lake to use as Eurasian Watermilfoil control.

The filtering process and bank stabilization that buffers provide help improve a lake's water quality, including water clarity. Studies in Minnesota, Maine and Michigan have shown that waterfront property value increases for every foot the water clarity of a lake increases. (Krysel et al, 2003).



Figure 14: Example of Inadequate Vegetative Buffer

Figure 15: Example of Adequate Buffer



Natural shoreland buffers serve important cultural functions. They enhance the lake's aesthetics. Studies have shown that aesthetics rank high as one of the reasons people visit or live on lakes. Shore buffers can provide visual & audio privacy screens for homeowners from other neighbors and/or lake users.

Adequate buffers on Sherwood Lake in some places could be easily installed on the inadequate areas by either letting the first 35 feet landward from the water just grow without mowing it, except for a path to the water, or by planting native seedlings sufficient to fill in the first 35 feet or using biologists to protect the shore that are vegetated. Where areas are deeply eroded, shaping, revegetating and protecting the shores will be necessary to prevent further erosion.



**Figure 16:
Heavily-eroded
Shore on
Sherwood Lake
that would need
installation of
shoreland
protection
practices**

WATER QUALITY

Between 2004 and 2006, Adams County Land & Water Conservation Department gathered water chemistry and other water quality information on Sherwood Lake. Historic information about water testing on Sherwood Lake was also obtained from the studies discussed earlier in this report.

Phosphorus

Most lakes in Wisconsin, including Sherwood Lake, are phosphorus-limited lakes: of the pollutants that end up in the lake, the one that most affects the overall quality of the lake water is phosphorus. The amount of phosphorus especially affects the frequency and density of aquatic vegetation and the frequency and density of various kinds of algae, as well as water clarity and other quality aspects. One pound of phosphorus can produce as much as 500 pounds of algae.

Phosphorus is not an element that occurs in high concentration naturally, so any lake that has significant phosphorus readings must have gotten that phosphorus from outside the lake or from internal loading. Some phosphorus is deposited onto the lake from atmospheric deposition, especially from soil or other particles in the air carrying phosphorus. A lake that includes a flooded wetland area may have a significant amount of phosphorus being released during the flushing of the wetland area. Phosphorus may accumulate in sediments from dying animals, dying aquatic plants and dying algae. If the bottom of the lake becomes anoxic (oxygen-depleted), chemical reactions may cause phosphorus to be released to the water column.

Although there are several forms of phosphorus in water, the total phosphorus (TP) concentration is considered a good indicator of a lake's nutrient status, since the TP concentration tends to be more stable than other types of phosphorus concentration. For an impoundment lake like Sherwood Lake, a total phosphorus concentration below 30 micrograms/liter tends to prevent nuisance algal blooms. Sherwood Lake's growing season (June-September) surface average total phosphorus level of 37.7 micrograms/liter is slightly over to the level at which nuisance algal blooms can be expected. And areas of Sherwood Lake do have nuisance-level algal blooms, especially in the shallower back bays.

Since phosphorus is usually the limited factor, measuring the phosphorus in a lake system thus provides an indication of the nutrient level in a lake. Increased phosphorus in a lake will feed algal blooms and also may cause excess plant growth.

The 2004-2006 summer average phosphorus concentration in Sherwood Lake places Sherwood Lake in the “fair” water quality section for impoundments, and in the “mesotrophic” level for phosphorus. The total epilimnetic phosphorus level has been creeping up in Sherwood Lake. In 1991, average epilimnetic total phosphorus was 24.33 micrograms/liter. By 2005, it averaged 35.6 micrograms/liter. The most recent figures, from summer 2007, show an average epilimnetic phosphorus level of 54.86 micrograms/liter, although the overall average for 2004-2007 is 41.99 micrograms/liter. These levels show that nutrients are accumulating in the lake as time goes on.






Figure 18: Average Eplimnetic Phosphorus Levels



As the Figure 18 indicates, the average growing season total phosphorus levels have varied and often registered above the level recommended to avoid nuisance algal blooms. However, the epilimnetic total phosphorus levels have stayed below the state impoundment average of 65 micrograms/liter. Especially due to the increasing epilimnetic total phosphorus levels, phosphorus should continue to be monitored and steps should be taken to reduce the phosphorus levels in the lake.

Groundwater testing of various wells around Sherwood Lake was done by Adams County LWCD and included a test one year for total phosphorus levels in the groundwater coming into the lake. The average TP level in the wells was 22.75 micrograms/liter, considerably lower than the lake surface water results.

Land use plays a major role in phosphorus loading. A key component of the computer models used is the phosphorus budget, that is, the estimated amount of phosphorus delivered to the lake from each land use type annually. The land uses that contribute the most phosphorus are non-irrigated agriculture and residences. Using the current land use data, as well as phosphorus readings from 2004 through 2006 water sampling, a phosphorus loading prediction model was run for Sherwood Lake. The current results are shown in Figure 19a.

Figure 19a: Current Phosphorus Loading by Land Use

MOST LIKELY ANNUAL PHOSPHORUS LOADING--Current		
	% Loading	Lbs/acre/yr
Grassland/Pasture	0.7%	8.03
Residential	5.2%	59.78
Water	1.7%	18.74
Woodland	2.0%	22.31
Septics	5.2%	58.89
Greater Surface Watershed	85.2%	917.52
Total		1085.27

In the prior studies, phosphorus loading was somewhat different than the current estimates (see figure 19b).

Figure 19b: Prior Phosphorus Loading Estimates

Land Use Phosphorus Contributions in lbs/acre/yr	1991	current
Grassland/Pasture	0.00	8.03
Residential	47.29	59.78
Water	0.00	18.74
Woodland	83.87	22.31
Septics	0.00	58.89
Greater Surface Watershed	1571.16	917.52
Total in pounds/acre/year	1702.32	1085.27

These figures reveal that overall phosphorus loading in the watershed is now less than it was in the early-1990s. This may be due to the installation of agricultural runoff practices in the upper watershed. However, contribution percentages by land use have changed. For example, contribution (not including septic) from residential areas was nearly twice as much in 2004 than it was in 1991, while contribution from the greater watershed (which includes agriculture) has gone down nearly 10%.

Figure 19c: Contribution Percentage by Land Use

MOST LIKELY ANNUAL PHOSPHORUS LOADING--Current		
	% Loading	lb/yr
Grassland/Pasture	0.7%	19.8
Residential	5.2%	147.4
Other Water	0.5%	13.2
Woodland	2.0%	55
Septics	5.2%	145.2
Lake Surface	1.2%	33
Greater Surface Watershed	85.2%	2395.58
Total in pounds/year	100.0%	2809.18

Currently, the most phosphorus loading is coming from the greater watershed, including irrigated agriculture, non-irrigated agriculture and upstream septic systems. Although phosphorus deposits such as that from flooded wetlands or from atmospheric deposition cannot be controlled by humans, phosphorus loads from human activities such as agriculture, residential development and septic systems can be partly controlled by changes in human land use patterns. Practices such as agricultural buffers, nutrient management, shoreland buffer restoration; infiltrating stormwater runoff from roof tops, driveways and other impervious surfaces; using no phosphorus lawn fertilizers; and reducing phosphorus input to and properly managing septic systems will minimize phosphorus inputs into the lake. Circumstances such as increased impervious surface, lawns mowed to water's edge, disturbance of shore areas, improperly-functioning septic systems and removal of native vegetation can greatly increase the volume and content of runoff—and thus increase the volume of phosphorus entering the lake. Many of these practices can also increase the concentration of phosphorus entering the lake, by runoff or other methods of entry.

The models were run using not only the current known phosphorus readings in the lake, but also representing decreases or increases of human-controlled phosphorus input by 10%, 25%, and 50%. Just a 10% reduction of the human-impacted phosphorus would reduce the overall load by 276.3 pounds/year. This figure may not seem like much---until you calculate that one pound of phosphorus can result in up to 500 pounds of algae. A 10% reduction in these three areas could result in up to 138,150 pounds less of algae per year!

Figure 20: Impact of Increase/Decrease on P Loading

LAND USE	lb/yr	-10%	-25%	-50%
Grassland/Pasture	19.8	17.82	14.85	9.90
Residential	147.4	132.66	110.55	73.70
Other Water	13.2	13.20	13.20	13.20
Woodland	55	49.50	41.25	27.50
Septics	145.2	130.68	108.90	72.60
Lake Surface	33	33.00	33.00	33.00
Greater Surface Watershed	2395.58	2156.02	1796.69	1197.79
Total	2809.18	2532.88	2118.44	1427.69

Reducing the amount of input from the surface and ground watersheds results in less nutrient loading into the lake itself. Under the modeling predictions, reducing phosphorus inputs from human-based activities even 10% could improve Sherwood Lake water quality by up to 8.6 micrograms/liter. A 25% reduction could save up to 21.5 micrograms/liter and could reduce the overall epilimnetic growing season total phosphorus to around the 30 micrograms/liter level to avoid nuisance algal blooms. These predictions make it clear that reducing current phosphorus inputs to the lake are essential to improve, maintain and protect Sherwood Lake’s health for future generations.






Figure 22: Photo of a Lake in Algal Bloom

Water Clarity

Water clarity is a critical factor for plants. If plants don't get more than 2% of the surface illumination, they won't survive. Water clarity can be reduced by turbidity (suspended materials such as algae and silt) and dissolved organic chemicals that color or cloud the water. Water clarity is measured with a Secchi disk. Average summer Secchi disk clarity in Sherwood Lake in 2004-2006 was 4.36 feet. This is poor water clarity, putting Sherwood Lake into the "eutrophic" category for water clarity.




Figure 23b: Secch Disk 1990-1999





Figure 23c: Secchi Disk 2002-2007



Sherwood Lake has a considerable history of Secchi disk readings in a number of years. A look at the average Secchi depth for the growing season in each year since 1986, Secchi disk depth readings on Sherwood Lake have generally decreased over the years (see figure 24). The overall average depth for the fourteen years for which there are records is 5.0 feet.

Figure 24: Average Growing Season Secchi Averages





Figure 25: Photo of Testing Water Clarity with Secchi Disk

Chlorophyll a

Chlorophyll-a concentrations provide a measurement of the amount of algae in a lake's water. Algae are natural and essential in lakes, but high algal populations can increase water turbidity and reduce light available for plant growth, as well as result in unpleasing odor and appearance. Studies have shown that the amount of chlorophyll a in lake water depends greatly on the amount of algae present; therefore, chlorophyll-a levels are commonly used as a water quality indicator. The 2004-2006 growing season (June-September) average chlorophyll concentration in Sherwood Lake was 20.7 micrograms/liter. Such an algae concentration places Sherwood Lake at the "poor" level for chlorophyll a results.







Figure 27 shows that summer chlorophyll-a averages stayed fairly steady from 1991 through 2005, staying in the “fair” category. However, in 2006 and 2007, chlorophyll-a levels shot up substantially. These two years coincide with many days of very hot and still weather. Such weather tends to encourage algal growth. Continued monitoring of chlorophyll-a will need to occur to determine if these high levels were the result of weather or are part of an ongoing trend.



Dissolved Oxygen

Oxygen dissolved in the water is essential to all aerobic aquatic organisms. The oxygen in a lake comes from the atmosphere and from the process of photosynthesis. Aquatic plants and algae consume carbon dioxide and respire oxygen back into the lake water. The distribution of oxygen within a lake is affected by many factors, including water circulation, water stratification, winds or storms, air temperature; water temperature, nutrient availability, and the density and location of algae and/or aquatic plants.

Prior studies of Sherwood Lake have found anoxic (no oxygen) or hypoxic (low oxygen) in Sherwood near the dam (also the location of the deepest water in Sherwood Lake) and in the south “arm” of the lake. However, during the 2004-2006 study, no anoxia was found and hypoxia of just of 5 milligrams/liter occurred only in June 2005. Generally, dissolved oxygen levels didn’t usually go below levels 5 milligrams/liter, the appropriate level for good fish survival. The charts (Figures 28) show the annual variations in dissolved oxygen levels in milligrams/liter, depth in feet and months of the years.




Figure 28a: Dissolved Oxygen Levels 2004 in milligrams/liter






Figure 28b: Dissolved Oxygen Levels During 2005 Water Testing in milligrams/liter











Figure 26c:
Dissolved Oxygen
Levels During
2006 Water Testing
in milligrams/liter

-  4/06
-  6/06
-  7/06
-  8/06
-  9/06

In deeper lakes, when the surface waters have cooled in autumn and water density throughout the water column is the same, the water column mixes vertically, a process known as “fall turnover.” Most of Sherwood Lake is shallow and does not stratify. However, the west end of the lake, where depths exceed 22 feet deep, does stratify and turns over in the spring or fall.

Further, since flowing stream goes through the lake from east to west through Sherwood Lake, some open water is common throughout the winter on part of the lake. This probably allows oxygen levels to stay elevated—even the winter, most of the dissolved oxygen readings over the amount needed by fish (over 5 milligrams/liter).

Figure 29a: One of the abundant fish in Sherwood Lake—Bluegill (*Lepomis macrochirus*)




Figure 29b: One of the common Fish in Sherwood Lake—Largemouth Bass (*Micropterus salmoides*)


Water Hardness, Alkalinity and pH

Testing done by Adams County LWCD on Sherwood Lake included annual testing for water alkalinity and water hardness. Hardness and alkalinity levels in a lake are affected by the soil minerals, bedrock type in the watershed, and frequency of contact between lake water & these materials.

Level of Hardness	Milligrams/liter CaCO ₃
SOFT	0-60
MODERATELY HARD	61-120
HARD	121-180
VERY HARD	>180

**Figure 30:
Hardness
Table**

One method of evaluating hardness is to test the water for the amount of calcium carbonate (CaCO₃) it contains. The surface water of all of the public access lakes in Adams County have water that is moderately hard to very hard, whether they are impoundments (man-made lakes) or natural lakes. In 2005 and 2006, random samples were also taken of wells around Sherwood Lake to measure the hardness of the water coming into the lake through groundwater. Hardness in the groundwater ranged from 180 (hard) to 208 (very hard), with an average of 187.2 milligrams/liter. Surface water hardness averaged 178 milligrams/liter, slightly lower than the groundwater hardness. The hardness in both surface and groundwater is likely due to the underlying bedrock in Adams County, which is mostly sandstone with pockets of dolomite and shale.



As the graph (Figure 31) shows, Sherwood Lake surface water testing results showed “hard” water (average 178 milligrams/liter CaCO₃), slightly more than the overall hardness average impoundments in Adams County of 166 milligrams/liter of Calcium Carbonate. Hard water lakes tend to produce more fish and aquatic plants than soft water lakes because they are often located in watersheds with soils that load phosphorus into the lake water.

Alkalinity is important in a lake to buffer the effects of acidification from the atmosphere. “Acid rain” has long been a problem with lakes that had low alkalinity level and high potential sources of acid deposition.


Acid Rain Sensitivity	ueq/l CaCO ₃
High	0-39
Moderate	49-199
Low	200-499
Not Sensitive	>500

Figure 32: Acid Rain Sensitivity

Well water alkalinity testing results ranged from 132 milliequivalents/liter to 180 milliequivalents/liter in alkalinity, with an average of 152 milliequivalents/liter. This is higher than the surface water average of 125.6 milliequivalents/liter. Sherwood Lake’s potential sensitivity to acid rain is moderate, but luckily for Adams County, the acid deposition rate is very low, probably due to the little industrialization in the county.

Alkalinity also affects the pH level of lake water. The acidity level of a lake’s water regulates the solubility of many minerals. A pH level of 7 is neutral. The pH level in Wisconsin lakes ranges from 4.5 in acid bog lakes to 8.4 in hard water, marl lakes.

Some of the minerals that become available under low pH, especially the metals aluminum, zinc and mercury, can inhibit fish reproduction and/or survival. Even what seems like a small variance in pH can have large effects because the pH scale is set up so that every 1.0 unit change increases acidity tenfold, i.e., water with a pH of 7 is 10 times more acid than water with pH of 8. Mercury and aluminum are not only toxic to many kinds of wildlife; they can also be toxic to humans, especially those that eat tainted fish.




The testing occurring from 2004-2006 also included regular monitoring of the pH at several depths in Sherwood Lake. As is common in the lakes in Adams County, Sherwood Lake has pH levels starting at just under neutral (6.88) at over 22 feet depth and increasing in alkalinity as the depth gets less. By 15 feet deep, the pH had risen to 7.47 and at 5 feet below the surface, the pH reached 7.53. A lake's pH level is important for the release of potentially harmful substances and also affects plant growth, fish reproduction and survival. Most plants grow best at pH levels between 5.5 and 8.

More importantly for many lakes, fish reproduction and survival are very sensitive to pH levels. The chart below indicates the effect of pH levels under 6.5 on fish (Figure 34):

Figure 34: Effects of pH Levels on Fish

Water pH	Effects
6.5	walleye spawning inhibited
5.8	lake trout spawning inhibited
5.5	smallmouth bass disappear
5.2	walleye & lake trout disappear
5	spawning inhibited in most fish
4.7	Northern pike, sucker, bullhead, pumpkinseed, sunfish & rock bass disappear
4.5	perch spawning inhibited
3.5	perch disappear
3	toxic to all fish


No pH levels taken in Sherwood Lake between 2004-2006 fell below the pH level that inhibits walleye reproduction. A lake with a neutral or slightly alkaline pH like Sherwood Lake is a good lake for fish and plant survival. Natural rainfall in Wisconsin averages a pH of 5.6. This means that if the rain falls on a lake without sufficient alkalinity to buffer that acid water coming in by rainfall, the lake's fish cannot reproduce. That is not a problem at Sherwood Lake. Sherwood Lake has a good pH level for fish reproduction and survival.



Other Water Quality Testing Results

CHLORIDE: Chloride does not affect plant and algae growth and is not known to be harmful to humans. It isn't common in most Wisconsin soils and rocks, so is usually found only in very low levels in Wisconsin lakes. However, the presence of a significant amount of chloride over a period of time indicates there may be negative human impacts on the water quality present from septic system failure, the presence of fertilizer and/or waste, deposition of road-salt, and other nutrients. An increased chloride level is thus an indication that too many nutrients are entering the lake, although the level has to be evaluated compared to the natural background data for chloride. The average chloride level found in Sherwood Lake during the 2004-2006 testing period was 11.73 milligrams/liter, elevated substantially above the natural level of 3 milligrams/liter for chloride in this area of Wisconsin.

Prior studies also found elevated chloride levels in Sherwood Lake. In fact, substantially elevated chloride levels have been found at Sherwood Lake since records were kept (1985). Further investigation as to the causes of such continued chloride elevations needs to be performed.



NITROGEN: Nitrogen is necessary for plant and algae growth. A lake receives nitrogen in various forms, including nitrate, nitrite, organic, and ammonium. In Wisconsin, the amount of nitrogen in a lake's water often corresponds to the local land use. Although some nitrogen will enter a lake through rainfall from the atmosphere, that coming from land use tends to be in higher concentrations in larger amounts,


coming from fertilizers, animal and human wastes, decomposing organic matter, and surface runoff. For example, the growth level of the exotic aquatic plant, Eurasian Watermilfoil (*Myriophyllum spicatum*) has been correlated with fertilization of lake sediment by nitrogen-rich spring runoff.

Nitrogen levels can affect other aspects of water quality. The sum of water testing results for nitrate, nitrite and ammonium levels of over .3 milligrams/liter in the spring can be used to project the likelihood of an algal bloom in the summer (assuming sufficient phosphorus is also present). Sherwood Lake combination spring levels from 2004 to 2006 averaged .99 milligrams/liter, above the .3 milligrams/liter predictive level for nitrogen-related algal blooms. These elevations suggest that some of the algal blooms on Sherwood Lake may be at least partly nitrogen-related.

CALCIUM and MAGNESIUM: Calcium is required by all higher plants and some microscopic lifeforms. Magnesium is needed by chlorophyllic plants and by algae, fungi and bacteria. Both calcium and magnesium are important contributors to the hardness of a lake's waters. Magnesium elevated about 125 milligrams/liter may have a laxative effect on some humans. Otherwise, no health hazards to humans and wildlife are known from calcium and magnesium. The average Calcium level in Sherwood Lake's water during the testing period was 41.36 milligrams/liter. The average Magnesium level was 17.7 milligrams/liter. Both of these are low-level readings.


SODIUM AND POTASSIUM: These elements occur naturally only in low levels in Wisconsin waters and soils. Their presence may indicate human-caused pollution. Sodium is found with chloride in many road salts and fertilizers and is also found in human and animal waste. Potassium is found in many fertilizers and also found in animal waste. The level of these two is generally not useful as a specific pollution indicator, but increasing levels or one or both of these elements can indicate possible contamination from damaging pollutants. High levels of sodium have also been found to influence the development of a large population of cyanobacteria, some of which can be toxic to animals and humans. Some health professionals have suggested that sodium levels over 20 milligrams/liter may be harmful to heart and kidney patients if ingested. From 2004-2006, the average sodium level was 3.4 milligrams/liter. Prior sodium levels, taken in the mid-1980's, were slightly lower at 2.1 milligrams/liter. The sodium level average from 1985 to 2006 was 2.14 milligrams/liter. The average potassium reading from 2004 to 2006 was 2.55 milligrams/liter. This is slightly higher than the potassium levels from the mid-1980s, which averaged 1.5 milligrams/liter. However, both of these reading remain low on the overall scale of potential problems.

SULFATE: In low-oxygen waters (hypoxic), sulfate can combine with hydrogen and becomes the gas hydrogen sulfate, which smells like rotten eggs and is toxic to most aquatic organisms. Sulfate levels can also affect the metal ions in the lake, especially iron and mercury, by binding them up, thus removing them from the water column. To prevent the formation of hydrogen sulfate, levels of 10 milligrams/liter are best. A health advisory kicks in at 30 milligrams/liter. Sherwood Lake sulfate levels averaged 29.01 milligrams/liter during the testing period, above the level for hydrogen sulfate formation, but still slightly below the health advisory level. However, the overall average for the years in which sulfate testing was done is 30.84, above the health advisory level.



TURBIDITY: Turbidity reflects water clarity. The term refers to suspended solids in the water column—solids that may include clay, silt, sand, plankton, waste, sewage and other pollutants. Turbid water may mask the presence of bacteria or other pollutants because the water looks murky or muddy. In general, turbidity readings of less than 5 NTU are best. Very turbid waters may not only smell, but also tend to be aesthetically displeasing, thus curtailing recreational uses of the water. Turbidity levels for Sherwood Lake’s waters were low in the 1980s, but have risen substantially since, with one reading over the 5 TU mark.

Figure 38: Turbidity Levels in Sherwood Lake



**Figure 39:
Examples of Very
Turbid Water**



HYDROLOGIC BUDGET

According to data in a 1971 WDNR bathymetric (depth) map, Sherwood Lake has 246 surface acres, and the volume of the lake is 2516 acre-feet. At that time, 7.5% of the lake was less than 3 feet deep and 8% was over 20 feet deep. The maximum depth was 27 feet.

A “hydrologic budget” is an accounting of the inflow to, outflow from and storage in a hydrological unit (such as a lake). “Residence time” is the average length of time particular water stays within a lake before leaving it. This can range from several days to years, depending on the type of lake, amount of rainfall, and other factors. “Flushing rate” is the time it takes a lake’s volume to be replaced. “Annual runoff volume”, as used in WiLMS, is the total water yield from the drainage area reaching the lake. The “drainage area” is the amount of area (in acres) contributing surface water runoff and nutrients to the lake. The “areal water load” is the total annual flow volume reaching the lake divided by the surface area of the lake. “Hydraulic loading” is the total annual volume of all water sources (including precipitation, non-point sources & point sources) loading into the lake.

Using the data gathered from historical testing and that done by the Adams County LWCD from 2004-2006, the WiLMS model calculated the tributary drainage area for Sherwood Lake as 2337.7 acres. The average unit runoff for Adams County in the Sherwood Lake area is 9.4 inches. WiLMS determined the expected annual runoff volume as 1831.2 acre-feet/year. Anticipated annual hydraulic loading is 5840.8 acre-feet/year. Areal water load is 23.7 feet/year. In an impoundment lake like Sherwood Lake, a significant portion of the water and its nutrient load running through it from the impounded creek tend to flush through the lake and continue downstream—in Sherwood Lake’s case, modeling estimates a water residence of 0.43 year. The calculated lake flushing rate is 2.32 1/year. Water and its load flow through Sherwood Lake fairly quickly.

Figure 40: Example of Hydrologic Budget





Figure 41: Sherwood Lake Bathymetric Map



TROPHIC STATE

The trophic state of a lake is one measure of water quality, basically defining the lake’s biological production status (see Figure 42). **Eutrophic lakes** are very productive, with high nutrient levels, frequent algal blooms and/or abundant aquatic plant growth. **Oligotrophic lakes** are those low in nutrients with limited plant growth and small populations of fish. **Mesotrophic lakes** are those in between, i.e., those which have increased production over oligotrophic lakes, but less than eutrophic lakes; those with more biomass than oligotrophic lakes, but less than eutrophic lakes; often with a more varied fishery than either the eutrophic or oligotrophic lakes. In comparing water quality testing results with the prediction from the computer modeling of this modeling with the actual figures outlined above, the actual Trophic State of Sherwood Lake is what was predicted from the modeling. Modeling results predicted that the overall TSI for Sherwood Lake would be **57**. This score places Sherwood Lake’s overall TSI at above average for impoundment lakes in Adams County (52.83).

Figure 42: Trophic Status Table

Score	<u>TSI Level Description</u>
30-40	<u>Oligotrophic:</u> clear, deep water; possible oxygen depletion in lower depths; few aquatic plants or algal blooms; low in nutrients; large game fish usual fishery
40-50	<u>Mesotrophic:</u> moderately clear water; mixed fishery, esp. panfish; moderate aquatic plant growth and occasional algal blooms; may have low oxygen levels near bottom in summer
50-60	<u>Mildly Eutrophic:</u> decreased water clarity; anoxic near bottom; may have heavy algal bloom and plant growth; high in nutrients; shallow eutrophic lakes may have winterkill of fish; rough fish common
60-70	<u>Eutrophic:</u> dominated by blue-green algae; algae scums common; prolific aquatic plant growth; high nutrient levels; rough fish common; susceptible to oxygen depletion and winter fishkill
70-80	<u>Hypereutrophic:</u> heavy algal blooms through most of summer; dense aquatic plant growth; poor water clarity; high nutrient levels

Sherwood Lake = 57


→

Phosphorus concentration, chlorophyll-a concentration and water clarity data are collected and combined to determine a trophic state. As discussed earlier, the average growing season epilimnetic total phosphorus for Sherwood Lake was 37.7 micrograms/liter. The average growing season chlorophyll-a concentration was 20.7 micrograms/liter. Growing season water clarity averaged a depth of 4.36 feet. Figure 39 shows where each of these measurements from Sherwood Lake falls in trophic level.


Figure 43: Sherwood Lake Trophic Status Overview

Trophic State	Quality Index	Phosphorus (ug/l)	Chlorophyll a (ug/l)	Secchi Disk (ft)
Oligotrophic	Excellent	<1	<1	>19
	Very Good	1 to 10	1 to 5	8 to 19
Mesotrophic	Good	10 to 30	5 to 10	6 to 8
	Fair	30 to 50	10 to 15	5 to 6
Eutrophic	Poor	50 to 150	15 to 30	3 to 4
Sherwood Lake		37.7	20.7	4.36

These figures show that Sherwood Lake has poor to fair levels overall for the three parameters often used to describe water quality: Secchi disk depths; average TP for the growing season; and chlorophyll a levels. It is normal for all of these values to fluctuate during a growing season. However, they can be affected by human use of the lake, by summer temperature variations, by algae growth & turbidity, and by rain or wind events. Sherwood Lake is above the county impoundments average for overall TSI levels—which is negative, since with TSI levels, the lower the better.



During the 1991 study, water chemistry samples were taken and TSI levels were calculated. Figure 45, which compares the current TSI levels to those from 1991, shows that all the parameters for TSI calculations have increased, indicating that the lake nutrient levels have risen substantially.




IN-LAKE HABITAT

Aquatic Plants

A healthy aquatic plant community plays a vital role within the lake community. This is due to the role plants play in improving water quality, providing valuable habitat resources for fish and wildlife, resisting invasions of non-native species and checking excessive growth of the most tolerant species.

An aquatic plant survey was performed in 2006 by staff from the Adams County Land & Water Conservation Department and a Tri-Lakes property owner. The aquatic plant community in Sherwood Lake is characterized by below average quality for Wisconsin lakes, poor species diversity and impacted by high levels of disturbance. Sherwood Lake is within the 25% of lakes in the state most tolerant of disturbance and furthest from an undisturbed condition. Disturbances include invasions of exotic species, boat traffic, shoreline development, harvesting and past herbicide treatments.

Of the 29 species found in Sherwood Lake, 25 were native and 4 were exotic invasives. In the native plant category, 14 were emergent, 1 was a free-floating plant, and 10 were submergent species. Four exotic invasives, *Myriophyllum spicatum* (Eurasian Watermilfoil), *Nasturtium microphyllum* (watercress), *Phalaris arundinacea* (Reed Canarygrass), and *Potamogeton crispus* (Curly-Leaf Pondweed) were found. Filamentous algae were found at 22.76% of the sample sites in 2006 and at 38.33% of the sites in 2000.



Of the plants on this list, several are species likely to increase in frequency and/or density in the case of regular drawdowns: *Lemna minor* (free-floating); *Najas flexilis* (submergent); *Potamogeton crispus* (submergent exotic); *Potamogeton pectinatus* (submergent); *Scirpus validus* (emergent) and *Potamogeton zosteriformis* (submergent). Some also tend to decrease with regular drawdowns: *Chara* spp (submergent); *Myriophyllum sibiricum* (submergent); *Myriophyllum spicatum* (submergent exotic); and *Vallisneria americana* (submergent). In general, regular drawdowns will tend to encourage the increase of plants that can survive frequent disturbances and will also tend to reduce the diversity of the aquatic plant community.

Figure 47. Sherwood Lake Aquatic Plant Species, 2006

<u>Scientific Name</u>	<u>Common Name</u>	<u>Type</u>	<u>Found in 2000</u>
<i>Calamagrostis canadensis</i>	Blue-Joint Grass	Emergent	
<i>Carex crawfordii</i>	Crawford's Sedge	Emergent	
<i>Carex comosa</i>	Longhair Sedge	Emergent	
<i>Ceratophyllum demersum</i>	Coontail	Submergent	x
<i>Chara spp</i>	Muskgrass	Submergent	x
<i>Eupatorium purpureum</i>	Sweetscented Joe Pye Weed	Emergent	
<i>Hypericum canadense</i>	Large St. John's Wort	Emergent	
<i>Iris versicolor</i>	Blue-Flag Iris	Emergent	
<i>Juncus spp</i>	Rush	Emergent	x
<i>Lathyris palustris</i>	Marsh Pea	Emergent	
<i>Lemna minor</i>	Small Duckweed	Free-Floating	x
<i>Myriophyllum sibiricum</i>	Northern Milfoil	Submergent	x
<i>Myriophyllum spicatum</i>	Eurasian Watermilfoil	Submergent	x
<i>Najas flexilis</i>	Bushy Pondweed	Submergent	x
<i>Nasturtium microphyllum</i>	Watercress	Floating-Leaf	
<i>Phalaris arundinacea</i>	Reed Canarygrass	Emergent	
<i>Potamogeton crispus</i>	Curly-Leaf Pondweed	Submergent	x
<i>Potamogeton illinoensis</i>	Illinois Pondweed	Submergent	
<i>Potamogeton pectinatus</i>	Sago Pondweed	Submergent	x
<i>Potamogeton pusillus</i>	Small Pondweed	Submergent	x
<i>Potamogeton zosteriformis</i>	Flat-Stem Pondweed	Submergent	x
<i>Renaunculus recurvatus</i>	Hooked Buttercup	Emergent	
<i>Sagittaria spp</i>	Arrowhead	Emergent	x
<i>Salix spp</i>	Willow	Emergent	
<i>Scirpus validus</i>	Soft-Stem Bulrush	Emergent	
<i>Thelypteris palustris</i>	Marsh Fern	Emergent	
<i>Typha latifolia</i>	Wide-Leaf Cattail	Emergent	x
<i>Vallisneria americana</i>	Water Celery	Submergent	x
<i>Zosterella dubia</i>	Water Stargrass	Submergent	x

Comparing the species found in 2006 to those reported in 2000, some changes are evident. Several plants found in 2006 were not found in 2000, especially emergents: *Carex crawfordii* (emergent); *Carex comosa* (emergent); *Eupatorium purpureum* (emergent); *Hypericum canadense* (emergent); *Iris versicolor* (emergent); *Lathyris palustris* (emergent); *Potamogeton illinoensis* (submergent); *Ranunculus recurvatus* (emergent); *Salix* spp (emergent); and *Scirpus validus* (emergent).

Potamogeton pectinatus, an aquatic plant favored by drawdowns, was the most frequently-occurring plant in Sherwood Lake in 2006. In 2000, the most frequent species was *Chara* spp. No species but *Potamogeton pectinatus* reached a frequency of 50% or greater in the lake overall in 2006, although *Chara* spp and *Potamogeton crispus* were not far under 50%, with occurrence frequencies of 45.53% and 42.28% respectively. In 2000, no aquatic species reached an overall occurrence frequency of over 50%.

Potamogeton pectinatus was also the densest plant in 2006 in Sherwood Lake. In the lake overall, none of the aquatic plant species had a mean density of over 2.0, meaning none occurred at more than average, in 2006. In 2006, the only species occurring at more than average density in any of the depth zones was *Potamogeton pectinatus* in the second (1.5 feet-5 feet) and third (5 feet-10 feet) depth zones. Densest in Depth Zone 1 (0 to 1.5 feet) was *Chara* spp; densest in the other three zones was *Potamogeton pectinatus*. No species occurred at more than average density in the lake overall in 2000, either. The only depth zone with more than average density of growth was Depth Zone 3, where *Chara* spp grew at more than average density.

However, when looking at the “mean density where present”, three plants in addition to *Potamogeton pectinatus* had a more than average “density where present” in 2006: *Chara* spp; *Eupatorium purpureum*; and *Potamogeton crispus*. This is lower than the seven species beside *Chara* spp that had more than average “density where present” in 2000: *Elodea canadensis*; *Juncus* spp; *Myriophyllum sibiricum*; *Nitella* spp; *Potamogeton foliosus*; *Potamogeton pusillus* and *Vallisneria americana*. These figures indicate some species in the lake have higher than average density that can interfere with fish habitat and recreational use.

Relative frequency and relative density are combined into a dominance value that demonstrates how dominant a species is within its aquatic plant community. Based on dominance value, *Chara* spp was the dominant aquatic “plant” species in Sherwood Lake in 2000. Sub-dominant was *Elodea canadensis*. However, in 2006, *Potamogeton pectinatus* dominated the aquatic plant community, with *Potamogeton crispus* and *Chara* spp next most dominant. The exotics found Sherwood Lake, were not present

in high frequency, high density or high dominance in either year although *Myriophyllum spicatum* had a greater frequency in 2000.

In 2006, *Potamogeton pectinatus* was dominant in Depth Zones 1 and 2, with *Chara* spp subdominant in each. *Potamogeton pectinatus* also dominated in Depth Zone 3, with *Potamogeton crispus* and *Ceratophyllum demersum* subdominant. *Potamogeton pectinatus* dominated Depth Zone 4 in 2006. In 2000, *Chara* spp dominated all four depth zones

Aquatic plants occurred at 84.6% of the sample sites in Sherwood Lake to a maximum rooting depth of 14 feet. This increased coverage from the 79.2% figure of 2000, when the maximum rooting depth was 12 feet. Filamentous algae were found in the three shallowest zones in both 2006 and 2000.

In 2006, the 1.5 feet-5 feet depth zone (Zone 2) produced the highest total occurrence of plant growth, followed closely by Depth Zone 3. There was then a slight drop in occurrence to Zone 1, then a sharp drop to Zone 4. The pattern was slightly different in 2000: Depth Zone 3 had the highest total occurrence, then a drop in frequency in Depth Zone 2. Depth Zone 1 was lower than Depth Zone 2, with Zone 4 having the lowest total occurrence of all. For plant density in 2006, Depth Zone 2 had the greatest total density, with Depth Zone 3 having slightly less. A sharp drop in density characterized Depth Zone 1 and even lower to Depth Zone 4. In 2000, the same pattern was followed.

Species richness increased slightly between 2000 and 2006, with the biggest increase in richness found in Depth Zone 1 (0-1.5 feet).

	2006	2000
Zone 1	5.38	2.35
Zone 2	2.94	2.58
Zone 3	2.8	3.56
Zone 4	1.95	1.69
Overall	2.98	2.93

Figure 48: Aquatic Plant Species Richness Table

The Simpson's Diversity Index for Sherwood Lake in 2006 was .84, indicating poor species diversity. A rating of 1.0 would mean that each plant in the lake was a different species (the most diversity achievable). This places it in the lowest quartile for Simpson's Diversity Index readings for both North Central Hardwood Forest and all Wisconsin lakes. This is lower than the Simpson's Diversity Index for 2000, which was .89. The 2006 AMCI for Sherwood Lake is 49, placing its quality below the

average for North Central Wisconsin Lakes and all Wisconsin Lakes. The AMCI value for 2000, 47, is also below average range for quality of aquatic plant community.

Figure 49: Aquatic Macrophyte Community Index-2006 & 2000

AMCI	2006	2006	2000	2000
Category	Result	Value	Result	Value
Max. Rooting Depth	14'	8	12'	6
% Littoral Zone Veg.	84.6%	10	79.2%	10
% Submersed Species	88%	9	95%	6
% Exotic Species	24%	3	8%	4
% Sensitive Species	4%	4	6%	5
Taxa #	23	9	16	8
Simpson's Index	0.84	6	0.89	8
		49		47

Using the AMCI index, some change has occurred in Sherwood Lake between 2000 and 2006, not necessarily for the better.

The presence of four invasive, exotic species could be a significant factor in the future. In 2006, *Potamogeton crispus* had the highest occurrence of any of the exotics found in Sherwood Lake, but *Myriophyllum spicatum* had an occurrence frequency of over 13% in 2006, despite the long history of both chemical and mechanical control efforts and despite this plant survey being done early in the summer, before some *Myriophyllum. spicatum* has reached its maximum growth. These plants must continue to be monitored, since their tenacity and ability to spread to large areas fairly quickly could make them a danger to the already low diversity of Sherwood Lake's current aquatic plant community.

The Average Coefficient of Conservatism and Floristic Quality Index calculation were performed on the field results. Technically, the Average Coefficient of Conservatism measures the community's sensitivity to disturbance, while the Floristic Quality Index measures the community's closeness to an undisturbed condition. Indirectly, they measure past and/or current disturbance to the particular community.

Previously, a value was assigned to all plants known in Wisconsin to categorize their probability of occurring in an undisturbed habitat. This value is called the plant's Coefficient of Conservatism. A score of 0 indicates a native or alien opportunistic invasive plant. Plants with a value of 1 to 3 are widespread native plants. Values of 4 to 6 describe native plants found most commonly in early successional ecosystem. Plants scoring 6 to 8 are native plants found in stable climax conditions. Finally,

plants with a value of 9 or 10 are native plants found in areas of high quality and are often endangered or threatened. In other words, the lower the numerical value a plant has, the more likely it is to be found in disturbed areas.

The Average Coefficient of Conservatism in Sherwood Lake in 2006 was 4.6, up slightly from 4.00 in 2000. This puts this lake in the lowest quartile for Wisconsin Lakes (average 6.0) and for lakes in the North Central Hardwood Region (average 5.6). The aquatic plant community in Sherwood Lake is in the category of those very tolerant of disturbance, probably due to selection by a series of past disturbances and heavy shoreline development.

The Floristic Quality Index of the aquatic plant community in Sherwood Lake of 16.85 in 2006 is in the lowest quartile for Wisconsin Lakes and the North Central Hardwood Region. This suggests that the plant community in Sherwood Lake is within the group of lakes farthest from an undisturbed condition in Wisconsin overall and in the North Central Hardwood Region. The 2000 figure of 16.97 was also in the lowest quartile. The Floristic Quality Index has decreased slightly between 2000 and 2006, suggesting more disturbance progress to the lake. Using either the Average Coefficient of Conservatism or the Floristic Quality Index scales, the aquatic plant community in Sherwood Lake apparently has been impacted by a high amount of disturbance.

“Disturbance” is a term that covers many disruptions to a natural community. It includes physical disturbances to plant beds such as boat traffic, plant harvesting, chemical treatments, dock and other structure placements, shoreline development and fluctuating water levels. Indirect disturbances like sedimentation, erosion, increased algal growth, and other water quality impacts will also negatively affect an aquatic plant community. Biological disturbances such as the introduction of non-native and/or invasive species (such as the Eurasian Watermilfoil, Reed Canarygrass and Curly-Leaf Pondweed found here), destruction of plant beds, or changes in aquatic wildlife can also negatively impact an aquatic plant community. Shore development and sediment deposition can also reduce the quality of the aquatic plant community.

Out of the 36 transects sampled on Sherwood Lake, only one site was entirely naturally vegetated. Therefore, no statistical evaluation comparing the aquatic macrophyte communities at disturbed vs. natural shores was appropriate.

Based on water clarity, chlorophyll and phosphorus data, Sherwood Lake is a eutrophic/mesotrophic impoundment with poor water clarity and fair to poor water quality. This trophic state should support substantial plant growth and several algal blooms.

Sufficient nutrients (trophic state), shallow lake, and nutrient input from heavy shore development on Sherwood Lake favor plant growth. Despite the sometime limiting effect of poor water clarity and sand sediments on aquatic plant growth, over 84% of the lake is vegetated, suggesting that even the heavily-sandy sediments in Sherwood Lake hold sufficient nutrients to maintain aquatic plant growth.

The very few shore areas of natural vegetation and wetlands on the lake should be preserved as they are to maintain habitat and to serve as a buffer for that area. Studies have suggested that runoff from naturally-buffered land is substantially less than that of developed areas. There are also some areas of deep erosion on steep banks that need to be addressed to prevent tree fall (and related root ball removal from bank) and bank preservation. Shoreline restoration of native vegetation is badly needed on Sherwood Lake and has actually decreased since 2000.

Some type of native vegetated shoreline covered only 21.46% of the lake shoreline in 2006, down from 30% in 2000. Disturbed shorelines—including bare sand, traditional cultivated lawn, hard structure (piers, decks, seawalls, etc.) and rock riprap--were the most frequently-occurring shore, covering 78.54% of the shore of Sherwood Lake in 2006, up from 70% of the shore in 2000.

Looking at the results from the 2000 survey and those from 2006 shows some changes in the aquatic plant community. There were more species found in 2006, and the structure of the aquatic plant community has changed with more emergent species present, but only one free-floating plant. No floating-leaf plants, which provide habitat and cover for fish and invertebrates, were found in either year. Further, when calculating the coefficient of similarity between the 2000 and 2006 surveys, they score as statistically dissimilar both in terms of frequency of occurrence and relative frequency. Based on frequency of occurrence, the aquatic plant communities of the two years are just over 45% similar. Using relative frequency, the score is only 53% similar. Similarity percentages of 75% or more are considered statistically similar. Obviously, the figures for Sherwood Lake are far below that figure.

It is worth noting that the report on the 2000 aquatic plant surveys mentioned the low level of emergent plants in Sherwood Lake. The 2006 survey shows that emergent plants were still scarce in Sherwood Lake, but there were more increased coverage from emergent plants in 2006.

Figure 50: Changes in the Aquatic Plant Community 2000 to 2006

	Changes in the Macrophyte Community			
	2000	2006	Change	%Change
Sherwood Lake	2000	2006	Change	%Change
Number of Species	18	25	7	38.89%
Maximum Rooting Depth	12.0	14.0	2	16.67%
% of Littoral Zone Unvegetated	20.80%	15.40%	-0.054	-25.96%
%Emergents	5.26%	12.50%	0.1	137.64%
%Free-floating	2.11%	0.00%	0.0	-100.00%
%Submergents	100.00%	100.00%	0.0	0.00%
%Floating-leaf	0.00%	0.00%	0.0	0.00%
Simpson's Diversity Index	0.89	0.84	-0.05	-5.51%
Species Richness	3.72	2.94	-0.78	-20.97%
Floristic Quality Index	16.97	16.85	-0.12	-0.71%
Average Coefficient of Conservatism	4	4.6	0.60	15.00%
AMCI Index	47	49	2.00	4.26%

The following aquatic species decreased between 2000 and 2006: *Ceratophyllum demersum*, *Chara* spp., *Myriophyllum spicatum*, *Najas flexilis*, *Potamogeton pusillus*, *Zosterella dubia*, *Sagittaria* spp, *Eleocharis acicularis*, *Nitella* spp., *Potamogeton foliosus*, and *Spirodela polyrhiza*. Species that increased included: *Myriophyllum spicatum* (an exotic invasive); *Potamogeton crispus* (an exotic invasive); *Potamogeton pectinatus*, *Potamogeton zosteriformis*; and *Vallisneria americana*. New species found in 2006 were mostly emergents, although two new invasives were also found: *Phalaris arundinacea* and *Nasturtium microphyllum*.

Figure 51a: Distribution of Emergent Plants in Sherwood Lake




Figure 51b: Distribution of Submergent Plants in Sherwood Lake



There was a long history of chemical use for treating aquatic plant growth and algae in Sherwood Lake, 1970-2000. Some chemical products that are now banned because of their toxicity were used. Broad-spectrum chemicals were also used. Two chemicals that do not biodegrade, but build up in the sediment, resulting in toxic sediment were used. .

Figure 52: Chemical Aquatic Plant Treatments in Sherwood Lake

Year	Copper (lbs)	Cutrine (gal)	Aquathol (gal)	Hydrothol (gal)	Diagquat (gal)	Rodeo (gal)	2,4-D (gal)	Silvex	AV-70
1970	250		10		5				
1971	305		17		14			3	
1972	293		9		20.5				
1973	620				12				
1974	1220				12		22		
1975	620		8.9		6.6		2		
1976	600		9.5		26				
1977	910		215	100	6				30
1978				550	8				8
1979	400								
1980	60			855					
1981	60			1200					
1982	450								
1983	500								
1984	200		27	1					
1985	70		56		8				
1986	900		38		6				
1987	430								
1988	605				6				
1989	50		7		5.5				
1990	400		20		22.5				
1991	200		3.5		18				
1992	250		10		8				
1993		15	9.5		10.5				
1994	360		17.5		10.5				
1995	425		13.25		5.25				
1996		32	14			14			
1997		72.5	2.5		2.5				
1999			6		6				
2000			35		35				
total	10178	119.5	528.65	2706	253.85	14	24	3	38

The problems with the herbicides that were used included the following:

- 1) One toxic compound used in Sherwood Lake was Silvex (2,2,4,5-TP). Silvex is now banned as a possible carcinogen.
- 2) The broad-spectrum chemical Diquat was used. This compound killed all plant species and inadvertently opened up areas for the introduction of exotic and invasive species. Over 253 gal of Diquat compounds were used over the 30-year span.
- 3) The Hydrothol formulation is toxic to young fish.
- 4) Cutrine and CuSO₄ are copper products that were used to kill algae and reduce swimmer's itch. Since copper is an element, it does not biodegrade further, building up the sediments. The drawbacks of copper treatments are: (a) the very short effective time; (b) the toxicity of copper to aquatic insects, an important part of the food chain in a lake; (c) the build up of copper in the sediments, resulting in sediments that are toxic to mollusks that are the natural consumers of algae in a lake.

Mechanical harvesting of aquatic plants in Sherwood Lake started in 1995 and has continued through 2006. The chart below shows the pounds of aquatic plant removed through mechanical harvesting through 2006. For 2005 and 2006, plant samples were taken to a laboratory to be tested for the amount of phosphorus in milligrams per kilogram of aquatic plants. This is also shown on the chart below.

Figure 53: Mechanical Harvesting on Sherwood Lake		
Year	Lbs Harvested	Phosphorus Removed (lbs)
1995	58,000	NA
1996	204,000	NA
1997	340,000	NA
1998	195,600	NA
1999	317,000	NA
2000	652,000	NA
2001	496,000	NA
2002	491,600	NA
2003	519,000	NA
2004	582,000	NA
2005	709,200	2147.81
2006	307,500	117.83
total	4,871,900	2,266

Ceratophyllum demersum
(Coontail)



Potamogeton pectinatus
(Sago Pondweed)

**Figure 54:
Some
Common
Native
Aquatic
Species in
Sherwood
Lake**



Chara spp
(Muskgrass)

Aquatic Plant Management Recommendations

- (1) Because the plant cover in the littoral zone of Sherwood Lake is at the top of the ideal (25%-85%) coverage for balanced fishery, continued harvesting to open fishing lanes could occur in some areas. Removal should occur by hand in the shallower areas to be sure that entire plants are removed and to minimize the amount of disturbance to the sediment.
- (2) Natural shoreline restoration and erosion control in many areas is needed, especially on some bare steep banks. If trees fall at the eroded sites due to continued erosion, large portions of the banks will fall with them. Natural shoreline has decreased since 2000 and disturbed shoreline has increased, especially in hard structure and rock riprap.
- (3) To protect water quality, a buffer area of native plants needs to be restored on those many sites that now have seawalls or have traditional lawns mowed to the water's edge. Most areas of the lake shoreline are unnatural and prone to erosion & runoff of nutrients & toxics. Unmowed native vegetation reduces runoff into the lake and filters runoff that enters the lake.
- (4) The Tri-Lakes Management District and the Sherwood Lake Association should continue to cooperate with the WDNR to monitor for zebra mussel introduction to protect the aquatic plant community in Sherwood Lake.
- (5) Studies indicate that properties around the lakeshore are putting nutrients into the lake, rather than most of the nutrients coming from the watershed. To improve the quality of the lake water and prevent further degradation:
 - (a) Stormwater management of the many impervious surfaces around the lake is essential to improve the quality of the lake water and prevent further degradation.
 - (b) No lawn chemicals should be used on properties around the lake. If they must be used, they should be used no closer than 50' to the shore. Green grass tends to equal green lake.
 - (c) The few sites where there is undisturbed shore should be maintained and left undisturbed.
 - (d) Sherwood Lake residents should identify, cooperate with and participate in watershed programs that will reduce nutrient and sediment inputs.
- (6) The aquatic plant management plan should be reviewed annually. Mechanical harvesting plans should continue target harvesting for Eurasian Watermilfoil

(EWM) and include target harvesting for Curly-Lead Pondweed to prevent further spread.

- (7) The Sherwood Lake Association may want to continue to apply for grants from the Wisconsin Department of Natural Resources to help defray the cost of aquatic plant management.
- (8) No broad-scale chemical treatments of aquatic plant growth are recommended due to the undesirable side-effects of such treatments, including increased nutrients from decaying plant material and decreased dissolved oxygen and opening up more areas to the invasion of EWM.
- (9) Any fallen trees should be left at the shoreline in the water for habitat.
- (10) The Tri-Lakes Management District conducted limited water quality monitoring for several years, but has decreased its involvement during 2004-2006 when Adams Land & Water Conservation Department was doing more intense monitoring as part of a Lake Classification Grant. Monitoring by the Lake District or through the DNR Self-Help Monitoring Program should be restarted.
- (11) No drawdowns of water level except for DNR-approved purposes should occur. Several of the plants found in Sherwood Lake in 2006 are those encouraged by drawdowns. In addition, water drawdowns are increasing the inflow of nutrient-rich groundwater into the lake.
- (12) The Tri-Lakes Management District should make sure that its lake management plan takes into account all inputs from both the Sherwood Lake surface ground watershed and inputs from Camelot & Sherwood Lakes, and addresses the concerns of this larger lake community.
- (13) Natural shoreline restoration should occur on disturbed sites.
 - (a) Unmowed native vegetation reduces shoreline erosion and run-off into the lake and filters the run-off that does enter the lake thus reducing nutrient inputs.
 - (b) Shoreline restoration could be as simple as leaving a band of natural vegetation around the shore by discontinuing mowing.
 - (c) Restoration could be as ambitious as extensive plantings of attractive native wetland species in the water and native grasses, flowers, shrubs and trees on the near shore area.

- (14) The Sherwood Property Owners Association & Tri-Lakes Management District should cooperate with programs in the watershed to reduce nutrient inputs to the lake. Currently nearly half of the relatively large watershed is in agriculture.

Aquatic Invasives

Sherwood Lake has five known invasive aquatic plant species: Curly-Leaf Pondweed (submergent); Eurasian Watermilfoil (submergent); Purple Loosestrife (emergent); Reed Canarygrass (emergent) and Watercress (floating-leaf). The lake gets a significant amount of transient boat traffic due to its location (right off a main highway) and large public boat ramp. The Tri-Lakes Management District has a lake management plan that includes management of aquatic invasives. The lake has been using targeted harvesting for Eurasian Watermilfoil, emphasizing the harvesting of that plant in May and September, while harvesting the summer months for navigation, rather than control of Eurasian Watermilfoil. In 2007, some lake citizens were trained to monitor the aquatic invasives and participate in the Clean Boats, Clean Waters boater education program.

Figure 55: Distribution of Exotic Aquatic Plants in Sherwood Lake



**Curly-Leaf
Pondweed**



**Figure 56: Invasive Aquatic Plants in
Known at Sherwood Lake**

Eurasian Watermilfoil



**Reed Canarygrass and
Purple Loosestrife**



Watercress

FISHERY/WILDLIFE/ENDANGERED RESOURCES


WDNR stocking and fishery inventories go back to 1968, when the lake was stocked after a chemical eradication of fish in 1967 to get rid of the rough fish population. Stocking in 1968 consisted of bluegills, largemouth bass, northern pike and walleye. A follow-up inventory in 1969 found that bluegills and pumpkinseeds were abundant; largemouth bass, northern pike, walleye and yellow perch were common; and shiners and white suckers were scarce. The most recent survey, done in 2002, found that bluegills and largemouth bass were abundant; black crappie, walleye and yellow perch were common; and northern pike was scarce. Between 1970 and 2000, thirteen other fish inventories were performed by the WDNR. In addition to those fish already mentioned, various fish surveys through the years also found brown bullheads, black bullheads, yellow bullheads, yellow suckers, golden shiners, and emerald shiners.

In 1999, the local WDNR fishery biologist reported that a recent survey of Sherwood Lake showed that the largemouth bass and northern pike populations were good and that those fish were healthy. However, although the panfish numbers were sufficient, he found them to be small and thin. He expressed concern about loss of invertebrate habitat that the fish fed on, about the chemicals killing zooplankton, and about the effects of the winter drawdown on the aquatic plant community.

Muskrat are also known to use Tri-Lakes shores for cover, reproduction and feeding. Seen during the field survey were various types of waterfowl and songbirds. Frogs and salamanders are known, using the lake shores for shelter/cover, nesting and feeding. Turtles and snakes also use this area for cover or shelter in this area, as well as nested and fed in this area. Upland wildlife feed and nest here as well.

There are several endangered resources in the Sherwood Lake surface watershed. Natural communities reported here include northern sedge meadow, northern wet forest, pine barrens and shrub-carr. Endangered plants known in the area include *Polygala cruciata* (crossleaf milkwort), *Juncus marginata* (grassleaf rush), and *Bartonia virginica* (yellow screwstem).

Figure 57: Photos of some of the species of concern in Sherwood Lake Watersheds*



YELLOW SCREWSTEM



CROSSLEAF MILKWORT

*information courtesy of Wisconsin Department of Natural Resources

RESOURCES

Bell, R.A. 2000. Phytoplankton Community Composition and Distribution in the Tri-Lakes Area. UW-Stevens Point.

Bryan, B., B. Charry. 2006. Conserving Wildlife in Maine's Shoreland Habitats. Maine Audobon Society.

Carlson, R.E. 1977. A Trophic State Index for Lakes. *Limnology and Oceanography* 22:361-369.

Comas, S.R. 1993. Tri-Lakes, Adams County, Wisconsin Lakes & Watershed Characterization. Blue Water Science, St. Paul, Minnesota.

Dennison, W., R. Orth, K. Moore, J. Stevenson, V. Carter, S. Kollar, P. Bergstrom, R. Batuik. 1993. Assessing Water Quality with Submersed Vegetation. *Bioscience* 43(2):86-94.

Engel, S. 1985. Aquatic Community Interactions of Submerged Macrophytes. Wisconsin Department of Natural Resources Bulletin #156.

Frankenberg, J. Land Use and Water Quality. Purdue Extension Publication ID-230.

James, T. 1992. A Guidebook for Lake Associations. The International Coalition for Land and Water Stewardship in the Red River Basin, Minnesota.

James, W.F., J. Barko & H. Eaken. 2002. Limnological Investigations of Camelot, Sherwood, & Arrowhead Lakes, WI. U.S. Army Corps of Engineers, Spring Valley, WI.

Kibler, D.F., ed. 1982. Urban Stormwater Hydrology. Water Resources Monograph 7. American Geophysical Union.

Krysel, C, E.M. Boyer, C. Parson, P. Welle. 2003. Lakeshore Property Values and Water Quality: Evidence from Property Sales in the Mississippi Headwaters Region. Report to the Legislative Commission on Minnesota Resources.

Lillie, R.A., J.W. Mason. 1983. Limnological Characteristics of Wisconsin Lakes. Department of Natural Resources Bulletin No. 138.

MSA Professional Services. 1999. Septic Study of Tri-Lakes.

Nichols, S. 1998. Floristic Quality Assessment of Wisconsin Lake Plant Communities with Example Applications. *Journal of Lake and Reservoir Management* 15(2):133-141.

Nichols, S., S. Weber, B. Shaw. 2000. A Proposed Aquatic Plant Community Biotic Index for Wisconsin Lakes. *Environmental Management* 26(5): 491-562.

Shaw, B., C. Sparacio, J. Stelzer, N. Turyk. 2001. Assessment of Shallow Groundwater Flow Chemistry & Interstitial Water Sediment, Aquatic Macrophyte Chemistry for Tri-Lakes, Adams County, WI. UW-Stevens Point.

Shaw, B., C. Mechanich, L. Klessing. Understanding Lake Data. UW-Extension Publication SR-02/2002-1M-525, 2000.

Terrell, C., P. Perfetti. 1989. Water Quality Indicators Guide: Surface Waters. United States Department of Agriculture Publication SCS-TP-161.

University of Wisconsin-Madison Water Resources Management Program. 1978. Fourteen Mile Creek Watershed: A Water Management Study. UW-Madison.

Wagner, C., J. Haack, R. Korth. Protecting Our Living Shores. 2003. Shoreland Stewardship Series #3 WDNR Publication WT-764-2003. UW-Extension, Wis. Lakes Partnership, WDNR, Wisconsin Association of Lake & River Alliance of Wisconsin.

Figure 58: Photo of Watercress on Sherwood Lake in Summer 2006



EXECUTIVE SUMMARY

An updated aquatic macrophytes (plants) transect field study of Sherwood Lake was conducted during July 2014 by staff from the Adams County Land and Water Conservatism Department and the Tri-Lakes Management District. The first quantitative vegetation study was performed by Wisconsin Department of Natural Resources staff in 2000, and follow-up surveys were performed in 2006 and 2009/2010 by the Adams County Land and Water Conservation Department and Tri-Lakes Management District staff.

Sherwood Lake, an impoundment of about 243 surface acres, is located in the Town of Rome, Adams County, Wisconsin. The dam impounds Fourteen-Mile Creek upstream from Arrowhead Lake and downstream from the two Camelot Lake dams, on its way to the Wisconsin River. There is a public boat launch on Sherwood Lake on the southwest edge of the lake owned by the Parks Department of Adams County with a small picnic area.

Of the forty-nine (49) species found in Sherwood Lake in 2015, forty-five (45) were native, and four were exotic invasives. In the native plant category, twenty-nine (29) species were emergent, three species were free-floating, and fourteen (13) were submergent aquatic species. There were no rooted floating-leaf plants found. Four exotic invasives, *Bromus inermis* (Smooth Brome), *Myriophyllum spicatum* (Eurasian Watermilfoil), *Phalaris arundinacea* (Reed Canarygrass), and *Potamogeton crispus* (Curly-Leaf Pondweed) were also found. The latter three have been found in previous aquatic plant surveys.

The most frequently occurring aquatic plant in the 2014 survey were Water Stargrass, found at over 46% of the sample sites, and Small Pondweed, found at over 44% of the 613 sample sites. No other 2014 species came close to this frequency of occurrence. The next most-frequently occurring species were Coontail, Northern Milfoil, and the invasive Curly-Leaf Pondweed. All five of these species are submergent plants.

Based on dominance value, Small Pondweed and Water Stargrass were nearly tied as the dominant aquatic plant species in Sherwood Lake in 2014. Small Pondweed alone was 14% of the aquatic plant community in 2014, with Water Stargrass following closely at 13.5%. No other species were close to these figures.

The Simpson's Diversity Index for Sherwood Lake in 2014 was .93, indicating very good species diversity. A rating of 1.0 would mean that each plant in the lake was a different species (the most diversity achievable). This places Sherwood Lake in the upper half for Simpson's Diversity Index readings for both North Central Hardwood Forest and all Wisconsin lakes. The Aquatic Plant Community Index (AMCI) value for 2014 was 59, down slightly from the 2011 score of 62. This score is at the top of the average AMCI range for all Wisconsin Lake and for the North Central Hardwood Forest Region.

MANAGEMENT RECOMMENDATIONS

1) Natural shoreline restoration and erosion control in several areas is still very much needed. The current Adams County Shoreland Ordinance will address these issues, since it requires buffers on all waterfront property by July 1, 2015. Despite

having several years to bring their properties into compliance, many Sherwood Lake property owners appear to have totally disregarded the ordinance and are running the risk of being fined. Cultivated lawn, mown right to the edge of the water, dominates much of the lake shore, rather than buffers. Buffers protect water quality and property values, among other things.

2) The Tri-Lakes Management District and the Sherwood Property Owner's Association should continue to cooperate with the WDNR to monitor for the presence of zebra mussels in the lake to protect the aquatic plant community.

3) Stormwater management of the many impervious surfaces around the lake is essential to maintain the current quality of the lake water and prevent further degradation. This is especially important because of the high amount of impervious surfaces around the lake.

4) No chemicals should be used on properties around the lake. If they must be used, they should be used no closer than 50 feet to the shore. This includes not just fertilizer, but also pesticides (sprayed or otherwise), and other contaminants.

5) Aquatic plant surveys should be repeated every 3 to 5 years to monitor changes in lake health and the aquatic plant community.

6) The lake management plan, including the aquatic plant management section, should continue to be reviewed annually.

7) Mechanical harvesting plans should not continue target harvesting for Eurasian Watermilfoil. Alternate methods of control need to be explored and implemented in addition to the current mechanical harvesting.

8) Continued evaluation of phosphorus removal by testing of the harvested plants should continue.

9) A plan for the management of curly-leaf pondweed needs to be developed and implemented.

10) The Tri-Lakes Management District may want to continue to apply for grants from the Wisconsin Department of Natural Resources to help defray the cost of aquatic plant management. It would also be a good idea to start setting aside some funds every year to build a reserve in case grants become unavailable.

11) No broad-scale chemical treatments of aquatic plant growth are recommended due to the undesirable side-effects of such treatments, including increased nutrients from decaying plant material and decreased dissolved oxygen and opening up more areas to the invasion of EWM.

12) Fallen trees should continue to be left at the shoreline or in the water to increase shore area habitat. Consultation with the WDNR Fisheries Biologist is recommended to evaluate the current habitat and gain recommendations for any improvement needed.

13) The Tri-Lakes Management District should continue involvement in water quality and invasive species monitoring through the Citizen Lake Monitoring

Program, the Clean Boats, Clean Waters program and grants for AIS management. Volunteers should be recruited and trained in case the funds for paying interns become unavailable.

14) Sherwood Lake residents should identify, cooperate with and participate in watershed programs that will reduce nutrient and sediment inputs. Nutrient levels appear to have increased within the lake watershed, so residents must take steps to reduce their nutrient inputs.

15) No drawdowns of water level except for DNR-approved purposes should occur. Drawdowns in the past have increased nutrient inputs to the lake through ground water; and these increased nutrients may be feeding the localized algae growth found in many places on Sherwood Lake. Some less desirable aquatic plant have decreased in frequency since 2006 and the cessation of annual drawdowns, and some more favorable aquatic plants have increased their presence since the winter drawdown stopped.

16) The few sites where there is undisturbed shore should be maintained and left undisturbed.

17) Cooperation with the Adams County Parks Department in keeping the boat ramp and small swimming area in safe condition should help reduce any negative impacts caused by the heavy use of these public areas.

18) A boat washing station at the park ramp area may help in decreasing other invasives from invading the lake. Portable boat washing stations can be installed

during the summer and fall months, then stored away during the winter months. This should decrease costs related to possible freezing pipes and winter vandalism.

19) The Tri-Lakes Management plan has long included having an on-the-ground survey of the watersheds, but this has not occurred. In 2013, such a survey was done on most of Big Roche a Cri Creek Watershed, the watershed just below 14-Mile Creek. That survey revealed that there were many problem areas of runoff, sloughing banks, and erosion. Since similar conditions occur in the 14-Mile-Creek Watershed, completing the survey is very important to identify problems areas and, hopefully, address them in the near future, as they may be contributing to water quality problems in the Tri-Lakes.

THE AQUATIC PLANT COMMUNITY FOR SHERWOOD LAKE, ADAMS COUNTY 2000-2014

I. INTRODUCTION

An updated aquatic macrophytes (plants) transect field study of Sherwood Lake was conducted during the summer of 2014 by the Adams County Land and Water Conservatism Department and the Tri-Lakes Management District. The first quantitative vegetation study was performed by Wisconsin Department of Natural Resources in 2000, and followup surveys were performed in 2006 and 2010/2011 by the Adams County Land and Water Conservation Department and Tri-Lakes Management District.

Information about the diversity, density, and distribution of aquatic plants is an essential component in understanding the lake ecosystem due to the integral ecological role of aquatic vegetation in the lake and the ability of vegetation to impact water quality (Dennison et al, 1993). This study will provide updated information useful for effective management of Sherwood Lake, including fish habitat improvement, protection of sensitive areas, aquatic plant management, and water resource regulation. This data will be compared to the past and used in future studies to offer insight into any changes in the lake.

Ecological Role: Lake plant life is the beginning of the lake's food chain, the foundation for all other lake life. Aquatic plants and algae provide food and oxygen for fish and wildlife, as well as cover and food for the invertebrates that many aquatic organisms depend on. Plants provide habitat and protective cover for aquatic animals. They also improve water quality, protect shorelines and lake bottoms, add to the aesthetic quality of the lake, and impact recreation.

Characterization of Water Quality: Aquatic plants can serve as indicators of water quality because of their sensitivity to water quality parameters such as clarity and nutrient levels (Dennison et al, 1993).

Testing has shown that Sherwood Lake has very hard water. The average reading was 180.4 milligrams/liter of Calcium Carbonate, placing Sherwood Lake on the hard to very hard line. Lake water pH has ranged from 5.92 to 8.46. Hard water lakes tend to produce more fish and aquatic plants than soft water lakes.

Background and History: Sherwood Lake, an impoundment of about 243 surface acres, is located in the Town of Rome, Adams County, Wisconsin. The dam impounds Fourteen-Mile Creek upstream from Arrowhead Lake and downstream from the two Camelot Lake dams, on its way to the Wisconsin River. Sherwood Lake is accessible off of State Highway 13 by turning east onto Queens Way.

There is a public boat launch on Sherwood Lake on the southwest edge of the lake operated by the Parks Department of Adams County. Heavy residential development around the lake is found along most of the lakeshore that includes many homes and several beach clubs. The surface watershed is 31.7% residential; 48.1% woodlands; 6.7% open grassland and 13.5% water. The ground watershed, which extends into Waushara County, has much irrigated and non-irrigated agriculture, except near to the lakes.

There are endangered or threatened resources in the watershed including the Karner Blue Butterfly, Grassleaf Rush, Yellow Screwstem, Crossleaf Milkwort; and the natural communities of northern sedge meadow, northern wet forest, pine

barrens, and shrub-carr. Archeological sites reported in the Sherwood Lake surface watershed include an unnamed burial site in the northern part of the watershed and the Bloody Nose Burial Mound south of the lake.

A fishery inventory in April 2002 revealed that yellow perch and largemouth bass are abundant in Sherwood Lake, while bluegills were common. Other fish, including walleye, black crappie, and northern pike, were scarce. Stocking records of the Wisconsin Department of Natural Resources go back to 1968. Stocking started after a chemical fish kill in 1967 to remove rough fish. Through the years, the lake has been stocked with largemouth bass, walleye, northern pike, and bluegills.

Soils in the Sherwood Lake surface watershed are sands of various slopes. These soils tend to be excessively-drained, with infiltration of water being rapid to very rapid, and permeability also high. They also usually have low water-holding and low organic matter content, thus making it difficult for vegetation establishment. These soils tend to be easily eroded by both water and wind.

Efforts at controlling aquatic plant growth have included both chemical treatments and mechanical harvesting. Once mechanical harvesting started in 1995, use of chemical decreased. 2000 was the last year the Tri-Lakes Management District used any chemicals for aquatic plant management. Both copper in pounds and cutrine in gallons added copper to Sherwood Lake. Copper is an element and does not degrade any further. Copper is known to harm native mollusks (clams, mussels, snails) and invertebrates that serve as food for the fish. Hydrothol, added to Sherwood Lake between 1977 and 1984, has been implicated in damage to young fish.

Mechanical harvesting is the major method of aquatic plant management in Sherwood Lake currently. Some hand-pulling of aquatic plants is used and lake residents are encouraged to hand-pull invasive aquatic plants.

FIGURE 1: SUMMARY OF MECHANICAL HARVESTING

<u>Year</u>	<u>Tons</u>	<u>Pounds</u>
1995	29.0	58,000
1996	102.0	204,000
1997	170.0	340,000
1998	97.8	195,600
1999	158.5	317,000
2000	326.0	652,000
2001	248.0	496,000
2002	245.8	491,600
2003	259.5	519,000
2004	291.0	582,000
2005	354.6	709,200
2006	153.7	307,400
2007	216.6	433,200
2008	252.3	504,600
2009	470.1	940,200
2010	374.7	749,400
2011	219.6	439,200
2012	446.9	893,700
2013	491.2	982,400
2014	432.9	876,000
TOTAL	5340.2	9,815,376

Harvested aquatic plant samples are sent to a laboratory to test the amount of phosphorus they contain. In 2014, over 158 pounds of phosphorus were removed from Sherwood Lake during the mechanical harvesting. Excess phosphorus is usually associated with the presence of nuisance-level aquatic plant growth and algae presence. Removal of phosphorus from the water column through harvesting helps reduce the total phosphorus levels in the lake.

Since the discovery of zebra mussels in Arrowhead Lake, the WDNR and citizen lake monitoring volunteers were monitoring Sherwood Lake for any sign of infestation. This has included veliger sampling and hanging mussel collection platforms at various areas throughout the lake. In 2009, Camelot Lake, upstream of Sherwood Lake on both ends to the east, was found to have zebra mussels. Adult zebra mussels were verified in Sherwood Lake in 2010 and continued to increase their presence afterwards. Many of the plants pulled from the lake in 2014 were covered with various-sized zebra mussels.

II. METHODS

Field Methods

The 2000, 2006, and 2009 studies were all based on the rake-sampling method developed by Jessen and Lound (1962), using stratified random transects. The shoreline was divided into 36 equal sections, with one transect placed randomly within each segment, perpendicular to the shoreline. The same transects were used for all three studies.

The 2011 and 2014 surveys were conducted using a modified Point Intercept Method (PI). This method involves calculating the surface area of a lake and dividing it (using a formula developed by the WDNR) into a grid of several points, always placed at the same interval from the next one(s). These points are related to a particular latitude and longitude reading. At each geographic point, the depth is noted and one rake is taken, with a score given between 1 and 3 to each species on the rake.

- A rating of 1 = a small amount present on the rake;
- A rating of 2 = moderate amount present on the rake;
- A rating of 3 = large amount present on the rake;

A visual inspection was done between points to record the presence of any species that didn't occur at the raking sites. Gleason and Cronquist (1991) nomenclature was used in recording plants found.

Data Analysis:

The percent frequency (number of sampling sites at which it occurred/total number of sampling sites) of each species was calculated. Relative frequency (number of species occurrences/total of all species occurrences) was also calculated. The mean density (sum of species' density rating/number of sampling sites) was calculated for each species. Relative density (sum of species' density/total plant density) was also calculated. "Mean density where present" (sum of species' density rating/number of sampling sites at which that species occurred) was calculated. Relative frequency and relative density results were summed to obtain a dominance value. Species diversity was measured by Simpson's Diversity Index.

The Average Coefficient of Conservatism and Floristic Quality Index were calculated as outlined by Nichols (1998) to measure plant community disturbance. A coefficient of Conservatism is an assigned value between 0 and 10 that measures the probability that the species will occur in an undisturbed habitat. The Average Coefficient of Conservatism is the mean of the coefficients for the species found in the lake. The Average Coefficient of Conservatism is used to calculate the Floristic Quality Index, a measure of a plant community's closeness to an undisturbed condition.

To measure the quality of the aquatic plant community, an Aquatic Macrophyte Index was determined using the method developed by Nichols et al (2000). This measurement looks at the following seven parameters and assigns each of them a number on a scale of 1-10: maximum depth of rooted plant growth; percentage of littoral zone vegetated; Simpson's diversity index; relative frequency of submersed species; relative frequency of sensitive species; taxa number; and relative frequency of exotic species. The average total for the North Central Hardwoods lakes and impoundments is between 48 and 57.

III. RESULTS


Physical Data

The aquatic plant community can be impacted by several physical parameters. Water quality, including nutrients, algae, and clarity, influence the plant community; the plant community in turn can modify these boundaries. Lake morphology, sediment composition, and shoreline use also affect the plant community.

The trophic state of a lake is a classification of water quality. Phosphorus concentration, chlorophyll-a concentration, and water clarity data are collected and combined to determine a trophic state. Eutrophic lakes are very productive, with high nutrient levels and large biomass presence. Oligotrophic lakes are those low in nutrients with limited plant growth and small fisheries. Mesotrophic lakes are those in between, i.e., those which have increased production over oligotrophic lakes, but less than eutrophic lakes; those with more biomass than oligotrophic lakes, but less than eutrophic lakes; those with a good and more varied fishery than either the eutrophic or oligotrophic lakes.

The limiting factor in most Wisconsin lakes, including Sherwood Lake, is phosphorus. Measuring the phosphorus in a lake system thus provides an indication of the nutrient level in a lake. Increased phosphorus in a lake will feed algal blooms and also may cause excess plant growth. Figures for total phosphorus during the growing season in Sherwood Lake go back to 1991. The average growing season total phosphorus level for Sherwood Lake from 1991 through 2014 is 35.6 micrograms/liter, above the recommended level for avoiding algal blooms (30 micrograms/liter), but below average for impoundments in Wisconsin (65 micrograms/liter). This concentration suggests that Sherwood Lake is likely to have some nuisance algal blooms. This places Sherwood Lake in the “fair” water quality section for impoundments, and in the “mesotrophic” level for total phosphorus.


Figure 2: Total Phosphorus Averages in Sherwood Lake



Chlorophyll-a is a pigment used by algae and aquatic plants in photosynthesis. Studies have shown that chlorophyll-a concentrations provide a measurement of the amount of algae in a lake’s water. Algae are natural and essential in lakes, but


high algal populations can increase water turbidity and reduce light available for plant growth. Chlorophyll-a records for Sherwood Lake are less complete than the total phosphorus information. The average growing season Chlorophyll-a level in Sherwood Lake from 2004 through 2014 was 20.4 micrograms/liter. These chlorophyll-a results place Sherwood Lake at the on the line between “mesotrophic” and “eutrophic”, suggesting “fair” to “poor” water quality in regards to chlorophyll-a.

Figure 3: Growing Season Averages for Chlorophyll-a



Water clarity is a critical factor for plants. If aquatic plants receive less than 2% of the surface illumination, they won't survive. Water clarity can be reduced by turbidity (suspended materials such as algae and silt) and dissolved organic chemicals that color or cloud the water. Water clarity is measured with a Secchi disk. Average summer Secchi disk clarity in Sherwood Lake in 1986 to 2014 was 5.3 feet. This is fair water clarity, putting Sherwood Lake into the “mesotrophic” category for water clarity.

Figure 4: Water Clarity Averages for Sherwood Lake



It is normal for all of these values to fluctuate during a growing season. They can be affected by human use of the lake, by summer temperature variations, by algae growth & turbidity, and by rain or wind events. Phosphorus tends to rise in early summer, then decline as late summer and fall progress. Chlorophyll- a tends to rise in level as the water warms, then decline as autumn cools the water. Water clarity also tends to decrease as summer progresses, probably due to algae growth, then declines as fall approaches.

These three parameters—water clarity, total phosphorus levels, and chlorophyll-a levels—are used to assign a lake the “trophic state” defined earlier in this report. Figure 5 shows where these averages for Sherwood Lake score.

FIGURE 5: TROPHIC STATE GRID

		TP (micro/l)	Chlo a (micro/l)	Secchi (feet)
Oligotrophic	Excellent	<1	<1	19
	Very Good	1 to 10	1 to 5	8 to 19
Mesotrophic	Good	10 to 30	5 to 10	6 to 8
	Fair	30 to 50	10 to 15	5 to 6
Eutrophic	Poor	50 to 150	15 to 30	3 to 4
	Very Poor	<150	<30	>3
Sherwood Lake		35.6	20.4	5.3

Lake Morphology

Lake morphology is an important factor in distribution of lake plants. Duarte & Kalff (1986) determined that the slope of a littoral zone could explain 72% of the observed variability in the growth of submerged plants. Gentle slopes support higher plant growth than steep slopes (Engel 1985). Sherwood Lake is a horseshoe-shaped narrow lake that lies in the middle of a series of lakes that are originally fed by a very large, multi-county stream system. Both Upper and Lower Camelot Lakes flow into Sherwood Lake. Much of the lake is shallow, although there are some areas of steeper drop-offs within the lake near the dam.

Most of the sediment in Sherwood Lake is hard, with little natural fertility and low available water holding capacity. Although such sediment may limit growth, most hard sediment sites in Sherwood Lake were vegetated. 81.9% of the sample sites were vegetated in Sherwood Lake, no matter what the sediment. Some of the unvegetated sites near shore appeared to have had aquatic vegetation cleared by hand harvesting.

Shoreline Land Use

Shoreline land use often strongly impacts the aquatic plant community and thus the entire aquatic community. Impacts can be caused by increased erosion and sedimentation and higher run-off of nutrients, fertilizers, and toxins applied to the land. Such impacts occur in both rural and residential settings. Despite some educational efforts by the Tri-Lakes Management District and the Adams County Land & Water Conservation Department, native vegetation in the shore areas has only increased by 4.4% since 2000. Bare sand and eroded areas have decreased slightly, probably replaced by the increase in herbaceous cover. Disturbed shorelines—including bare sand, traditional cultivated lawn, hard structure (piers, decks, seawalls), and rock riprap--have actually increased slightly overall, although cultivated lawn is down by about 5% since 2000. There continue to be an excess of cultivated lawn lots to the shore, with minimal buffer areas.

MACROPHYTE DATA

Of the forty-nine (49) species found in Sherwood Lake in 2015, forty-five (45) were native, and four were exotic invasives. In the native plant category, twenty-nine (29) species were emergent, three species were free-floating, and fourteen (13) were submergent aquatic species. There were no rooted floating-leaf plants found. Four exotic invasives, *Bromus inermis* (Smooth Brome), *Myriophyllum spicatum* (Eurasian Watermilfoil), *Phalaris arundinacea* (Reed Canarygrass), and *Potamogeton crispus* (Curly-Leaf Pondweed) were also found. The latter three have been found in previous aquatic plant surveys.

FIGURE 6: AQUATIC SPECIES FOUND IN 2014


Scientific Name	Common Name	Type
<i>Alisma triviale</i>	Northern Water Plantain	Emergent
<i>Amaranthus retroflexus</i>	Redroot	Emergent
<i>Asclepias incarnata</i>	Swamp Milkweed	Emergent
<i>Athyrium filix-femina</i>	Lady Fern	Emergent
<i>Bromus inermis</i>	Smooth Brome	Emergent
<i>Calamagrostis canadensis</i>	Blue-Joint Grass	Emergent
<i>Carex aquatilis</i>	Water Sedge	Emergent
<i>Carex comosa</i>	Bristly Sedge	Emergent
<i>Carex crawfordii</i>	Crawford's Sedge	Emergent
<i>Carex hystericina</i>	Bottlebrush Sedge	Emergent
<i>Ceratophyllum demersum</i>	Coontail	Submergent
<i>Chara contraria</i>	Opposite Stoneword	Submergent
<i>Cicuta bulbifera</i>	Bulb-Bearing Water Hemlock	Emergent
<i>Cornus spp</i>	Dogwood	Emergent
<i>Cyperus strigosus</i>	False Nut Sedge	Emergent
<i>Echinochloa walterii</i>	Coast Barnyard Grass	Emergent
<i>Elodea canadensis</i>	Common Waterweed	Submergent
<i>Impatiens capensis</i>	Jewelweed	Emergent
<i>Iris versicolor</i>	Blue-Flag Iris	Emergent
<i>Lactuca canadensis</i>	Tall Lettuce	Emergent
<i>Lemna minor</i>	Lesser Duckweed	Free-Floating
<i>Lycopus americanus</i>	American Bugleweed	Emergent
<i>Lycopus uniflorus</i>	Northern Bugleweed	Emergent
<i>Lysimachia terrestris</i>	Swamp Candle	Emergent
<i>Mentha arvensis</i>	Field Mint	Emergent
<i>Myriophyllum heterophyllum</i>	Various-Leaved Milfoil	Submergent
<i>Myriophyllum sibiricum</i>	Northern Milfoil	Submergent
<i>Myriophyllum spicatum</i>	Eurasian Watermilfoil	Submergent
<i>Najas flexilis</i>	Bushy Pondweed	Submergent
<i>Onoclea sensibilis</i>	Sensitive Fern	Emergent
<i>Phalaris arundinacea</i>	Reed Canarygrass	Emergent
<i>Potamogeton amplifolius</i>	Large-Leaf Pondweed	Submergent
<i>Potamogeton crispus</i>	Curly-Leaf Pondweed	Submergent
<i>Potamogeton foliosus</i>	Leafy Pondweed	Submergent
<i>Potamogeton friesii</i>	Fries' Pondweed	Submergent
<i>Potamogeton pusillus</i>	Small Pondweed	Submergent
<i>Potamogeton zosteriformis</i>	Flat-Stemmed Pondweed	Submergent
<i>Ranunculus aquatilis</i>	White Water Crowfoot	Submergent
<i>Rumex</i>	Water Dock	Emergent
<i>Sagittaria latifolia</i>	Common Arrowhead	Emergent
<i>Salix exigua</i>	Sandbar Willow	Emergent
<i>Schoenoplectus. tabernaemontani</i>	Soft-Stemmed Bulrush	Emergent
<i>Solanum dulcamara</i>	Bittersweet Nightshade	Emergent
<i>Sparganium emersum</i>	Narrow-Leaved Burreed	Emergent

<i>Spirodela polyrhiza</i>	Greater Duckweed	Free-Floating
<i>Stuckenia pectinata</i>	Sago Pondweed	Submergent
<i>Typha spp</i>	Cattail	Emergent
<i>Vallisneria americana</i>	Water Celery	Submergent
<i>Verbena hastata</i>	Blue Vervain	Emergent
<i>Wolffia columbiana</i>	Common Watermeal	Free-Floating
<i>Zosterella dubia</i>	Water Stargrass	Submergent

FREQUENCY OF OCCURRENCE

The most frequently occurring aquatic plants in the 2014 survey were Water Stargrass, found at over 46% of the sample sites, and Small Pondweed, found at over 44% of the 613 sample sites. No other 2014 species came close to these frequencies of occurrence. The next most-frequently occurring species were Coontail, Northern Milfoil, and the invasive Curly-Leaf Pondweed. All five of these species are submergent plants.

Figure 7: Most Frequently-Occurring Aquatic Plants 2014



DENSITY OF OCCURRENCE

None of the aquatic species found during the 2014 exhibited extra dense growth throughout the entire lake. There was some localized density. Most aquatic plants grow where the light can reach them. Secchi disc readings are used to predict maximum rooting depth for plants in a lake (Dunst, 1982). Following this formula, the average 2014 Secchi reading for Sherwood Lake was 5.8 feet. Using that figure, the formula would be $(5.8 \text{ feet} \times 1.22) + 2.73 \text{ ft} = 7.08 + 2.73 = 9.8 \text{ feet}$. This is in keeping with the depth zone in which the highest density of plants occurred: between 5 and 12 feet in depth. There were almost no “heavy” rakes where the density of plants found was marked ‘3’ on the scale explained earlier.

DOMINANCE

Relative frequency and relative density are combined into a dominance value that demonstrates how dominant a species is within its aquatic plant community. Based on dominance value, Small Pondweed and Water Stargrass were nearly tied as the dominant aquatic plant species in Sherwood Lake in 2014. Small Pondweed alone was 14% of the overall aquatic plant community in 2014, with Water Stargrass following closely at 13.5%. No other species were close to these figures.


Eurasian Watermilfoil (EWM) and Curly-Leaf Pondweed (CLP) have varied wildly in regards to their position in the aquatic plant community in Sherwood Lake. There are a number of potential explanations for these variations: timing of the survey, weather conditions of the spring and summer, level of boat traffic, and so on. However, these two invasives have steadily remained a part of the aquatic plant community in Sherwood Lake for at least 15 years. Because of their

potential to spread easily, continued monitoring of their presence level is necessary.

Figure 8: Percent of Overall Aquatic Plant Community of CLP & EWM

	2000	2006	2009	2011	2014
Curly-Leaf Pondweed	3.5%	17.0%	1.5%	6.0%	6.50%
Eurasian Watermilfoil	2.0%	4.5%	12.5%	7.0%	5.50%

Figure 9: Dominant Species in 2014




DISTRIBUTION

In 2014, aquatic plants occurred at 81.9% of the sample sites in Sherwood. This is similar to the overall vegetation level in Sherwood Lake since 2000. The distribution of the various types of aquatic plants is shown on the following maps.

As noted earlier, emergent plants continue to be rare in Sherwood Lake. Many of the shallow water areas in front of lakefront properties have large areas of cleared

lakebed. In addition, due to the large amount of cultivated lawn as well as significant areas of rock riprap, growth of nearshore emergent plants is not favored.


Figure 10: DISTRIBUTION OF EMERGENT & FREE-FLOATING AQUATIC PLANTS FOUND IN SHERWOOD LAKE IN 2014



Submergent plants dominate the lake, as they do most of the impoundments in Adams County. Of the four aquatic plant structure types—emergent, rooted floating-leaf, free-floating, and submergent—Sherwood Lake is totally lacking rooted free-floating plants (such as water lilies), has only a few free-floating plants

(located mostly in small bays), and has only a few emergents. The only place that submergents are not found is in the very deep areas near the dam structure.

Figure 11: SUBMERGENT PLANTS (in blue)



As is evident in the following maps, both Curly-Leaf Pondweed and Eurasian Watermilfoil are found throughout Sherwood Lake, except in the deeper areas near the dam. However, there are more of both in the lower arm of the lake, perhaps because that is the part of the lake most accessible from the public boat launch,

which is located at the southwestern curve of the lower arm of Sherwood Lake. This lower arm tends to get much more traffic, especially tubing, water-skiing and jet-skiing, than the upper arm of the lake.

Figure 12: Distribution of Curly-Leaf Pondweed 2014 (in lavender)





Figure 13: Distribution of Eurasian Watermilfoil 2014 (in red)



THE COMMUNITY

The Simpson's Diversity Index for Sherwood Lake in 2014 was .93, indicating very good species diversity. A rating of 1.0 would mean that each plant in the lake was a different species (the most diversity achievable). This places Sherwood Lake in the upper half for Simpson's Diversity Index readings for both North Central Hardwood Forest and all Wisconsin lakes. The Aquatic Plant Community Index (AMCI) value for 2014 was 59, down slightly from the 2011 score of 62. This score is at the top of the average AMCI range for all Wisconsin Lake and for the North Central Hardwood Forest Region.

Figure 14: 2014 Aquatic Macrophyte Community Index Values

2014	parameter	score
Maximum rooting depth	21	10
% of littoral zone vegetated	82.2	10
% of submergent plants (on relative frequency)	85	10
% of invasives (on relative frequency)	14	4
% of sensitive species (on relative frequency)	9	5
Simpson's Index of Diversity	0.93	10
Taxa # (number of species)	48	10
Total Score		59

The Average Coefficient of Conservatism and Floristic Quality Index calculation were performed on the field results. Technically, the Average Coefficient of Conservatism measures the community's sensitivity to disturbance, while the Floristic Quality Index measures the community's closeness to an undisturbed condition. Indirectly, they measure past and/or current disturbance to the particular community. Previously, a value was assigned to all plants known in Wisconsin to categorize their probability of occurring in an undisturbed habitat. This value is called the plant's Coefficient of Conservatism. A score of 0 indicates a native or alien opportunistic invasive plant. Plants with a value of 1 to 3 are widespread native plants. Values of 4 to 6 describe native plants found most commonly in early successional ecosystem. Plants scoring 6 to 8 are native plants found in stable climax conditions. Finally, plants with a value of 9 or 10 are native plants found in areas of high quality and are often endangered or threatened. In other words, the lower the numerical value a plant has, the more likely it is to be found in disturbed areas.

The Average Coefficient of Conservatism for Sherwood Lake in 2014 was 4.4. This figure has stayed between 4.0 and 5.0 since the first survey was done in 2000. This puts this lake in the lowest quartile for Wisconsin Lakes (average 6.0) and for

lakes in the North Central Hardwood Region (average 5.6). The aquatic plant community in Sherwood Lake is in the category of those very tolerant of disturbance, probably due to selection by a series of past disturbances and heavy shoreline development.

Figure 15: Floristic Quality and Coefficient of Conservatism of Sherwood Lake, Compared to Wisconsin Lakes and Northern Wisconsin Lakes.

	Average Coefficient of Conservatism †	Floristic Quality ‡
Wisconsin Lakes	5.5, 6.0, 6.9 *	16.9, 22.2, 27.5
NCHR	5.2, 5.6, 5.8 *	17.0, 20.9, 24.4
Sherwood Lake 2014	4.4	30.57

* - Values indicate the highest value of the lowest quartile, the mean and the lowest value of the upper quartile.

† - Average Coefficient of Conservatism for all Wisconsin lakes ranged from a low of 2.0 (the most disturbance tolerant) to a high of 9.5 (least disturbance tolerant).

‡ - lowest Floristic Quality was 3.0 (farthest from an undisturbed condition) and the high was 44.6 (closest to an undisturbed condition).

The Floristic Quality Index is a tool that can be used to identify areas of high conservation value, monitor sites over time, assess the anthropogenic (human-caused) impacts affecting an area and measure the ecological condition of an area (M. Bourdaghs, 2006). The Floristic Quality Index of the 2014 aquatic plant community in Sherwood Lake was 30.57. This puts Sherwood Lake in the upper quarter for all Wisconsin Lakes (16.9-27.5) and in the upper quartile for North Central Hardwood Lakes (median 17-24.4). Although this figure is higher than average, it has to be evaluated in the context of all the other results that can be used to look at the aquatic plant community in Sherwood Lake (water quality results, Average Coefficient of Conservatism, AMCI, etc.), it is clear that Sherwood

Lake's aquatic plant community has been impacted by a fair amount of disturbance.

“Disturbance” is a term that covers many disruptions to a natural community. It includes physical disturbances to plant beds such as boat traffic, plant harvesting, chemical treatments, dock and other structure placements, shoreline development, and fluctuating water levels. Indirect disturbances like sedimentation, erosion, increased algal growth, and other water quality impacts may also negatively affect an aquatic plant community. Biological disturbances such as the introduction of non-native and/or invasive species (such as the Eurasian Watermilfoil, Reed Canarygrass and Curly-Leaf Pondweed found here), destruction of plant beds, or changes in aquatic wildlife can also negatively impact an aquatic plant community. Shore development and sediment deposition can also reduce the quality of the aquatic plant community.

Disturbances likely to have impacted the aquatic plant community of Sherwood Lake include: substantial boat traffic and overall heavy recreational use; lack of shore buffers on most of the waterfront properties; presence of several aquatic invasives; internal and external nutrient loading; algae growth; erosion of and sedimentation from the soft sandy shores; the land use changes in the upper watershed.

IV. COMPARISONS TO PRIOR SURVEYS

The biggest changes in the aquatic plant community since 2006 have been in the increase in the occurrence frequency of emergent plants and decrease in the occurrence frequency of submergent plants. For more than 20 years, there was a

drawdown of Sherwood Lake every winter. These ceased about 2006-2008. There is an increase in the presence of emergents, but they seemed to have stabilized, still comprising a small amount of the overall aquatic plant community.

Figure 16: Plant Types Over the Years by Percentage of Community

	2000	2006	2011	2014
Emergent	2.5	7.5	7	7
Floating-Leaf	0	1	0	0
Free-Floating	2	0	1.5	1
Submergent	95.5	91.5	91.5	92


Another change has been the reduction of Sago Pondweed (*Stuckenia pectinata*), which once had an occurrence frequency of over 73%. This plant is encouraged by disturbances such as winter drawdowns. By 2009, its occurrence frequency had declined to just over 40% and by 2014, it was down to just over 18%.

As noted earlier, the two main aquatic invasives in Sherwood Lake since 2000 have remained Curly-Leaf Pondweed (CLP) and Eurasian Watermilfoil (EWM). Their presence in the overall aquatic plant community has varied greatly, with EWM reaching its highest presence in 2009, while CLP reached its high in 2006.

Since both the 2011 and 2014 surveys were conducted using the same survey method, the aquatic plant communities for those years were compared in regards to actual frequency of occurrence and relative frequency of occurrence, using the coefficient of similarity. This is an index, first developed by Jaccard in 1901, which compares the similarity and diversity of sample sets. In this instance, the figure considers the frequency of occurrence and relative frequency of all species found, then determines how similar the overall aquatic plant communities are.

Similarity percentages of 75% or more are considered statistically similar (Dennison et al, 1993).

Figure 17: CLP & EWM by Percentage of Aquatic Plant Community



According to the information gained from the 2011 and 2014 surveys, the aquatic plant communities in those two years were only 54.8% similar based on actual frequency of occurrence and 61.5% based on relative frequency. Both of these figures are below the 75% statistically similar range. However, these figures are between the prior comparisons between the 2006/2000 and 2009/2006 communities. For the 2006/2000 comparison, the results were 45.5% similar based on actual frequency of occurrence and 53.7% similar based on relative frequency. The comparison between the 2009/2006 communities were slightly better at 70.3% on actual occurrence frequency and 69.0% based on relative frequency. This suggests the aquatic plant community in Sherwood Lake has not stabilized yet, which may also make it especially vulnerable to colonization by new invasives.

Figure 18: Changes in the Macrophyte Community 2011-2014

Sherwood	2011	2014	Change
Number of Species	42	48	6
Maximum Rooting Depth	19.4	21.0	2
% of Littoral Zone Vegetated	69.3	82.2	12.9
%Sites/Emergents	7%	5%	0.0
%Sites/Free-floating	1%	1%	0.0
%Sites/Submergents	82%	84%	0.0
Simpson's Diversity Index	0.91	0.93	2.00
Species Richness	1.76	3.14	1.38
Floristic Quality	26.74	31.47	4.73
Average Coefficient of Conservatism	4.28	4.54	0.26
AMCI Index	60	59	-1.00

Figure 19 provides an example of some of the benefits provided to various birds, fish and mammals by some of the plants found in Sherwood Lake in 2009.

Figure 19: Uses of Some Aquatic Plants

	<u>Fish</u>	<u>Water</u>	<u>Shore</u>	<u>Upland</u>	<u>Muskrat</u>	<u>Beaver</u>	<u>Deer</u>
		<u>Fowl</u>	<u>Birds</u>	<u>Birds</u>			
<i>Ceratophyllum demersum</i>	F,I,C,S	F,I,C			F		
<i>Chara</i>	F,S	F,I,C					
<i>Lemna minor</i>	F,I,C,S	F	F		F	F	
<i>Myriophyllum heterophyllum</i>	F,I,C,S	F,I	F		F		
<i>Myriophyllum sibiricum</i>	F,I,C,S	F,I	F		F		
<i>Najas flexilis</i>	F,C	F	F				
<i>Stuckenia pectinata</i>	F,I,C,S	F,I	F		F	F	F
<i>Potamogeton zosteriformis</i>	F,I,C,S	F,I	F		F	F	F
<i>Schoenoplectus tabernaemontani</i>	F,C,I	F,C	F,C,N	F	F	F	F
<i>Typha spp</i>	I,C,S	F	F,C,N		F,C,N	F	
<i>Spirodela polyrhiza</i>	F,I,C,S	F	F		F	F	

F = Food; I = Shelters Invertebrates; C = Cover; S = Spawning; N = Nesting

V. DISCUSSION

Based on water clarity, chlorophyll-a, and phosphorus data, Sherwood Lake is a mesotrophic/mildly eutrophic impoundment lake with fair water clarity and poor to

fair water quality. This trophic state should support substantial plant growth and localized algal blooms.

Sufficient nutrients (trophic state), hard water, good water clarity, shallow lake, and nutrient-rich inputs from increased shore development at Sherwood Lake favor plant growth. Despite the sometime limiting effect of sand sediments on aquatic plant growth, most of the lake is vegetated, suggesting that even the sand sediments in Sherwood Lake hold sufficient nutrients to maintain aquatic plant growth.

Historically, many aquatic plant treatments in Sherwood Lake were chemical. There has been mechanical harvesting to try to reduce plant growth in the last 10 years. A continued regular schedule and pattern of machine harvesting will help in removing vegetation from the lake and may somewhat help with nutrient reduction. Any machine harvesting should be designed to avoid spreading Eurasian Watermilfoil. Aesthetically, it might also help to skim off the high density of filamentous algae and free-floating plants in certain areas of the lake.

Of the forty-nine (49) species found in Sherwood Lake in 2015, forty-five (45) were native, and four were exotic invasives. In the native plant category, twenty-nine (29) species were emergent, three species were free-floating, and fourteen (13) were submergent aquatic species. There were no rooted floating-leaf plants found. Four exotic invasives, *Bromus inermis* (Smooth Brome), *Myriophyllum spicatum* (Eurasian Watermilfoil), *Phalaris arundinacea* (Reed Canarygrass), and *Potamogeton crispus* (Curly- Leaf Pondweed) were also found. The latter three have been found in previous aquatic plant surveys.

There a few areas of native vegetation and wetland shores on the lake that should be preserved as they are to maintain habitat and to serve as a buffer for that area. Studies have suggested that runoff from such land is substantially less than that of developed areas. There are also some areas of deep erosion on steep banks that need to be addressed to prevent tree fall (and related root ball removal from bank) and bank preservation.

The Simpson’s Diversity Index for Sherwood Lake in 2014 was .93, indicating very good species diversity. This scores places Sherwood Lake in the upper range for all Wisconsin Lakes (.80 to .90) and for the North Central Hardwood Region (.82 to .90).

Figure 20: Changes in SI and AMCI Scores

	2000	2006	2009	2011	2014
SI	0.89	0.84	0.87	0.91	.93
AMCI	48	50	55	60	59

The 2014 AMCI for Sherwood Lake was 59, placing its quality in the upper range for North Central Hardwood Region Lakes (48 to 57) and all Wisconsin Lakes (45 to 57).

VI. CONCLUSION


Sherwood Lake is a mesotrophic to eutrophic impoundment with poor to fair water quality and water clarity. Structurally, the aquatic plant community contains emergent plants and a few submergent plants, but has no rooted floating-leaf and little free-floating aquatic vegetation.

A healthy and diverse aquatic plant community plays a vital role within the lake ecosystem. Plants help improve water quality by trapping nutrients, debris and pollutants in the water body; by absorbing and/or breaking down some pollutants; by reducing shore erosion by decreasing wave action and stabilizing shorelines and lake bottoms; and by tying-up nutrients that would otherwise be available for algae blooms. Aquatic plants provide valuable habitat resources for fish and wildlife, often being the base level for the multi-level food chain in the lake ecosystem, and also produce oxygen needed by animals.

Further, a healthy and diverse aquatic plant community can better resist the invasion of species (native and non-native) that might otherwise “take over” and create a lower quality aquatic plant community. A well-established and diverse plant community of natives can help check the growth of more tolerant (and less desirable) plants that would otherwise crowd out some of the more sensitive species, thus reducing diversity.

Vegetated lake bottoms support larger and more diverse invertebrate populations that in turn support larger and more diverse fish and wildlife populations (Engel, 1985). Also, a mixed stand of aquatic macrophytes (plants) supports 3 to 8 times more invertebrates and fish than do monocultural stands (Engel, 1990). A diverse plant community creates more microhabitats for the preferences of more species.

Figure 21: Aquatic Ecosystem Web



MANAGEMENT RECOMMENDATIONS

1) Natural shoreline restoration and erosion control in several areas is still very much needed. The current Adams County Shoreland Ordinance will address these issues, since it requires buffers on all waterfront property by July 1, 2015. Despite having several years to bring their properties into compliance, many Sherwood Lake property owners appear to have totally disregarded the ordinance and are running the risk of being fined. Cultivated lawn, mown right to the edge of the water, dominates much of the lake shore, rather than buffers. Buffers protect water quality and property values, among other things.

- 2) The Tri-Lakes Management District and the Sherwood Property Owner's Association should continue to cooperate with the WDNR to monitor for the presence of zebra mussels in the lake to protect the aquatic plant community.
- 3) Stormwater management of the many impervious surfaces around the lake is essential to maintain the current quality of the lake water and prevent further degradation. This is especially important because of the high amount of impervious surfaces around the lake.
- 4) No chemicals should be used on properties around the lake. If they must be used, they should be used no closer than 50 feet to the shore. This includes not just fertilizer, but also pesticides (sprayed or otherwise), and other contaminants.
- 5) Aquatic plant surveys should be repeated every 3 to 5 years to monitor changes in lake health and the aquatic plant community.
- 6) The lake management plan, including the aquatic plant management section, should continue to be reviewed annually.
- 7) Mechanical harvesting plans should not continue target harvesting for Eurasian Watermilfoil. Alternate methods of control need to be explored and implemented in addition to the current mechanical harvesting.
- 8) Continued evaluation of phosphorus removal by testing of the harvested plants should continue.

9) A plan for the management of curly-leaf pondweed needs to be developed and implemented.

10) The Tri-Lakes Management District may want to continue to apply for grants from the Wisconsin Department of Natural Resources to help defray the cost of aquatic plant management. It would also be a good idea to start setting aside some funds every year to build a reserve in case grants become unavailable.

11) No broad-scale chemical treatments of aquatic plant growth are recommended due to the undesirable side-effects of such treatments, including increased nutrients from decaying plant material and decreased dissolved oxygen and opening up more areas to the invasion of EWM.

12) Fallen trees should continue to be left at the shoreline or in the water to increase shore area habitat. Consultation with the WDNR Fisheries Biologist is recommended to evaluate the current habitat and gain recommendations for any improvement needed.

13) The Tri-Lakes Management District should continue involvement in water quality and invasive species monitoring through the Citizen Lake Monitoring Program, the Clean Boats, Clean Waters program and grants for AIS management. Volunteers should be recruited and trained in case the funds for paying interns become unavailable.

14) Sherwood Lake residents should identify, cooperate with and participate in watershed programs that will reduce nutrient and sediment inputs. Nutrient levels

appear to have increased within the lake watershed, so residents must take steps to reduce their nutrient inputs.

15) No drawdowns of water level except for DNR-approved purposes should occur. Drawdowns in the past have increased nutrient inputs to the lake through ground water; and these increased nutrients may be feeding the localized algae growth found in many places on Sherwood Lake. Some less desirable aquatic plant have decreased in frequency since 2006 and the cessation of annual drawdowns, and some more favorable aquatic plants have increased their presence since the winter drawdown stopped.

16) The few sites where there is undisturbed shore should be maintained and left undisturbed.

17) Cooperation with the Adams County Parks Department in keeping the boat ramp and small swimming area in safe condition should help reduce any negative impacts caused by the heavy use of these public areas.

18) A boat washing station at the park ramp area may help in decreasing other invasives from invading the lake. Portable boat washing stations can be installed during the summer and fall months, then stored away during the winter months. This should decrease costs related to possible freezing pipes and winter vandalism.

19) The Tri-Lakes Management plan has long included having an on-the-ground survey of the watersheds, but this has not occurred. In 2013, such a survey was done on most of Big Roche a Cri Creek Watershed, the watershed just below 14-Mile Creek. That survey revealed that there were many problem areas of runoff,

sloughing banks, and erosion. Since similar conditions occur in the 14-Mile-Creek Watershed, completing the survey is very important to identify problems areas and, hopefully, address them in the near future, as they may be contributing to water quality problems in the Tri-Lakes.

LITERATURE CITED

Bourdaghs, M., C.A. Johnston, and R.R. Regal. 2006. Priorities and performances of the floristic quality index in great lakes coastal wetlands. *Wetlands* 26(3):718-736.

Dennison, W., R. Orth, K. Moore, J. Stevenson, V. Carter, S. Kollar, P. Bergstrom and R. Batuik. 1993. Assessing water quality with submersed vegetation. *BioScience* 43(2):86-94.

Duarte, Carlos M. and Jacob Kalff. 1986. Littoral slope as a predictor of the maximum biomass of submerged macrophyte communities. *Limnol.Oceanogr.* 31(5):1072-1080.

Dunst, R.C. 1982. Sediment problems and lake restoration in Wisconsin. *Environmental International* 7:87-92.

Engel, Sandy. 1985. Aquatic community interactions of submerged macrophytes. Wisconsin Department of Natural Resources, Technical Bulletin #156. Madison, WI.

Evans, Reesa. 2006. The aquatic plant community of Sherwood Lake, Adams County, WI. 2000-2006. Adams County Land & Water Conservation Department.

Evans, Reesa. 2011. The aquatic plant community of Sherwood Lake, Adams County, WI. 2000-2011. Adams County Land & Water Conservation Department.

Gleason, H, and A. Cronquist. 1991. *Manual of Vascular Plants of Northeastern United States and Adjacent Canada* (2nd Edition). New York Botanical Gardens, N.Y.

Jaccard, P. 1901. Etude comparative de la distribution florale dans une poitve des Alpes et des Jura (in translation). Bulletin de la Socrete Vaudoise des Sciences Naturalles.

Jackson, H.O. and W.C. Starrett. 1959. Turbidity and sedimentation at Lake Chataqua, Illinois. Journal of Wildlife Management 14:157-168.

Jessen, Robert, and Richard Lound. 1962. An evaluation of a survey technique for submerged aquatic plants. Minnesota Department of Conservatism. Game Investigational Report No. 6.

Konkel, Deborah. 2000. The aquatic plant community of Sherwood Lake, Adams County, WI. Wisconsin Department of Natural Resources. Eau Claire.

Nichols, Stanley, and R.L. Nichols, ed. 1974. Mechanical and Habitat Manipulation for Aquatic Plant Management. Wisconsin Department of Natural Resources Technical Bulletin #77.

Nichols, Stanley. 1998. Floristic quality assessment of Wisconsin lake plant communities with example applications. Journal of Lake and Reservoir Management 15(2):133-141.

Nichols, S., S. Weber and B. Shaw. 2000. A proposed aquatic plant community biotic index for Wisconsin lakes. Environmental Management 26(5):491-502.

North Carolina State University Water Quality Group. Date Unknown. "Algae". Water Resource Characterization Series.

Quigley, M. March 1996. NOAA Public Affairs Bulletin 96-111.

Shaw, B., C. Mechenich and L. Klessig. 1993. Understanding Lake Data. University of Wisconsin-Extension. Madison, WI.

TRI-LAKES MANAGEMENT PLAN		
<u>AQUATIC SPECIES MANAGEMENT: GOALS and ACTIONS ITEMS</u>	<u>WHO</u>	<u>WHEN</u>
Machine harvest aquatic plants to improve water quality, to provide safe boating areas, and to improve aquatic habitat.		
1. Harvest maps will be followed by harvesting crew. Areas in the lake 6 feet deep or greater will be harvested as needed to a depth of 5 feet. Areas shallower than 6 feet deep can have a harvested path of 60 feet wide with a harvest depth of 4 feet or less. The path will get no closer to the shoreline than the docks. No machine harvesting in the areas between the shoreline and the ends of the boat docks. A 30-foot wide path may be hand-harvested in these areas by the property owner. Harvest period will be between mid-May and end of September.	Tri-Lakes Mgmt District	annually
2. Conservacy areas will be marked on the harvest maps.	Tri-Lakes Mgmt District	annually
Monitor the harvesting of aquatic plants.		
1. WDNR representative and a Tri-Lakes Management representative will meet annually to inspect the harvesting operations.	Tri-Lakes Mgmt District WDNR	annually
2. Record the pounds of aquatic plants removed by taking an average weight of a trailer-full of plants harvested and multiplying this by the number of trailer loads. This will be documented and reported to the WDNR Aquatic Plant Specialist by 12/31.	Tri-Lakes Mgmt District	annually
3. Composite samples will be taken randomly from the harvested plans (per protocol provided by the WDNR) and sent to a certified lab to measure the phosphorus content. This is done to determine the amount of phosphorus being removed from the lakes by harvesting plants.	Tri-Lakes Mgmt District WDNR	annually
4. Add GPS system to harvester to improve efficiency. Explore alternative product for marine application for better efficiency.	Tri-Lakes Mgmt District	ongoing
Eliminate the use of chemicals to control aquatic plants to improve water quality. This excludes noxious weed and invasive plant control.		
1. Educate individuals on aquatic plant control methods by placing article developed by Adams LWCD in Lake Association communications.	Lake Associations Adams LWCD	annually

Update 2006 Aquatic Plant Community Survey		
1. WDNR Lake Classification grant specifies aquatic plant survey to be conducted. Aquatic plants to be identified and quantified. Completed in 2006; Camelot completed 2009; Arrowhead completed in 2010. Sherwood completed in 2011. Camelot completed in 2013. Arrowhead & Sherwood completed 2014. Reports posted on Adams County LWCD website.	Adams County LWCD WDNR	2014
Prevent the spread of & try to eliminate invasive & noxious species.		
1. Invasive plants and noxious weeds will be mapped annually & in full aquatic plant surveys.	Adams County LWCD, WDNR	annually
2. Develop & implement plan to: control existing exotic, invasive & noxious species; and prevent the introduction of exotic, invasive and noxious species. This plan will include both educational & boat inspection components. CBCW program started in 2010 & has continued every summer since then.	Adams County LWCD WDNR Tri-Lakes Mgmt District private consultants	ongoing
3. Make annual contacts with boat dealers, boat patrol & water ski team & landscapers regarding methods to prevent spread of invasives.	Tri-Lakes Mgmt District	annually
4. Pursue financial assistance for implementation of prevention & control plan.	Adams County LWCD, WDNR Tri-Lakes Mgmt District Private Consultants	December 1 each year
5. Recontact Adams County Parks Committee to pursue installing boat washing equipment at public boat landings.	Adams LWCD & Parks Tri-Lakes Mgmt District	2016-2017
6. Control Purple Loosestrife using pulling, chemical spot treatments & bio-controls.	Adams LWCD, others	Ongoing
TRI-LAKES MANAGEMENT PLAN		
<u>AQUATIC SPECIES MANAGEMENT: GOALS and ACTIONS ITEMS (con'td)</u>		
	<u>WHO</u>	<u>WHEN</u>
Manage impacts to lakes resulting from the zebra mussel invasion.		
1. Arrange for a series of articles & provide material to the Wisconsin Rapids newspaper, Adams-Friendship newspaper, fishing clubs & lake associations.	Tri-Lakes Mgmt District	ongoing

2. Use volunteer group to provide education.	Tri-Lakes Mgmt District Lake Associations	ongoing
3. Develop an advertisement to be placed in lake association newsletters & local newspapers to notify the public that there are zebra mussels in Lakes Arrowhead & Camelot & Sherwood.	Tri-Lakes Mgmt District Adams LWCD, WDNR	ongoing
4. Seek funding from AIS grant for boat launch attendant through the Clean Boats, Clean Waters program.	Tri-Lakes Mgmt District Adams LWCD, WDNR	ongoing
Monitor the impact of zebra mussels.		
1. Monitor zebra mussel research on control methods.	Tri-Lakes Mgmt District Adams LWCD, WDNR	annually
2. Improve coordination and communications with WDNR Specialist & Adams LWCD.	Tri-Lakes Mgmt District Adams LWCD, WDNR	annually
<u>DAMS: GOALS and ACTIONS ITEMS</u>		
<u>WHO</u>		
<u>WHEN</u>		
Maintain and operate Arrowhead, Sherwood, and Camelot Dams to: insure public safety, proper dam function, and a stable lake level.		
1. Conduct inspections and record findings as specified in WDNR standards	engineer certified by Nat'l Assoc. of Prof. Eng.	per DNR schedule
2. Operate, inspect, and repair dams to meet Wisconsin laws in State Statute Chapter 31 and NR Chapter 330.	Adams LWCD	at least 3 times/wkly
3. Review and update the Emergency Action Plan.	Adams LWCD	annually
4. Implement the Emergency Action Plan in times of emergencies. Evacuation route signs added in 2014 by Adams County.	County Board Chair, LWCD Emer Mgmt, Town of Rome	as needed
5. Maintain water level gauges that are placed on the sides of the water flow control structures. These gauges will be the official tool used to determine lake levels. Lake level information may be obtained on line starting Juen 2010.	Adams LWCD	ongoing
6. Maintain history of levels; data can be obtained from Adams County LWCD at 608-339-4268.	Adams LWCD	annually

<u>RECREATION: GOALS and ACTION ITEMS</u>	<u>WHO</u>	<u>WHEN</u>
Manage lakes for boating and skiing.		
1. Provide areas for boating/skiing as part of the harvest plan.	Tri-Lakes Mgmt District	ongoing
2. Regulate lake traffic to increase lake safety. This will be accomplished by implementing & enforcing state laws, county and town ordinances. Pursue 2nd boat purchase & increase awareness with Town of Rome.	Town of Rome WDNR	ongoing
3. Implement methods of enforcing state laws, county and town ordinances. Methods may include anonymous tips hotline, goodwill ambassador groups, & education. Once methods have been identified, develop strategy to implement.	Tri-Lakes Mgmt District Town of Rome	ongoing
4. Install no-wake buoys 100 feet off the face of the dams shortly after ice-out.	Adams County LWCD	ongoing
Manage lakes for fishing.		
1. Work with WDNR Fisheries Biologist about stocking bass, panfish & forage fish in the lakes as funding allows.	Lake Associations Tri-Lakes Mgmt District	ongoing
2. Educate lake users about VHS & bait rules. Contact bait shops also.	Tri-Lakes Mgmt District Adams LWCD	ongoing
3. Educate lake users about improve littoral & riparian zones to improve fish habitat as long as navigation not impeded. Permit may be required.	Tri-Lakes Mgmt District	ongoing
Manage lakes for swimming.		
1. IMaintain safety of public swim area on Camelot Lake by marking with buoys	Town of Rome Adams Park & Rec.	ongoing
2. Report any illnesses from blue-green algae to www.dhs.wi.gov/eh/bluegreenalgae	Tri-Lakes Mgmt District	ongoing
Improve and develop parks associated with the Tri-Lakes		
1. Coordinate efforts to develop plan to improve and develop Tri-Lakes parks. Improvements include adding boat washing stations & containers to dispose of unused bait & aquatic plants at boat landings.	Town of Rome Adams Park & Rec. Lake Associations	ongoing

TRI-LAKES MANAGEMENT PLAN		
<u>NUTRIENT REDUCTION: GOALS and ACTION ITEMS</u>	<u>WHO</u>	<u>WHEN</u>
Reduce phosphorus and ammonia entering groundwater that then enters into Tri-Lakes.		
1. Inventory and map septic systems within the Tri-Lakes Management District. Manage system to notify owners (pre-7/1/1992) to pump septic systems.	Tri-Lakes Mgmt District	ongoing
2. Verify county & state-approved method of sanitary disposal on camping lots.	Town of Rome	ongoing
Reduce nutrients entering the Tri-Lakes by surface water.		
1. Continue upstream ditch testing started in 2007 and augmented in 2014, testing for phosphorus, nitrogen, chloride & total suspended solids, as well as flow amount. Use information to identify problem areas & develop practices to address input.	Adams LWCD Tri-Lakes Mgmt District	ongoing
2. Conduct shore habitat assessment around lakes & streams in watershed, using WDNR protocol.	Adams LWCD Tri-Lakes Mgmt District	2017-2018
3. Use information to develop plans to address identified problems areas.	Adams LWCD Tri-Lakes Mgmt District Lake Associations	2018-2019
4. Educate shoreline owners on the benefits of riparian buffers and storm water runoff management utilizing meetings, newsletters, and individual visits.	Tri-Lakes Mgmt District Adams LWCD, P & Z	ongoing
5. Plan/design & offer cost share to install shoreline protection, riparian buffers, and storm water runoff management to areas identified in inventory. Utilize Adams Soil & Water Resource Mgmt Program & WDNR financial assistance programs. Cost-sharing grants may be available to help.	Adams LWCD WDNR, landowners Lake associations	ongoing
6. Assist lakeshore property owners with Shoreland Protection Ordinance riparian compliance. Cost-sharing grants available to help. Compliance deadline is July 1, 2015.	Tri-Lakes Mgmt District Adams LWCD	ongoing
7. Monitor methods for sanitary disposal for year-round campers. Review town ordinance.	Town of Rome Adams County	ongoing
8. Develop compliance plan for shoreland protection buffer conformance, including plan for dealing with beach clubs.	Adams P & Z	2017-2018

<u>WATER QUALITY: GOALS and ACTION ITEMS</u>	<u>WHO</u>	<u>WHEN</u>
Reduce algae blooms and nutrient levels.		
1. Utilize the Citizen Monitoring Program to monitor water quality to measure water clarity, chlorophyll-a and total phosphorus.	Adams LWCD, Lake Assoc Tri-Lakes Mgmt District	ongoing
2. Monitor water quality at six sites in the upper watershed. Monitor flow rates at 3 sites	Adams LWCD Tri-Lakes Mgmt District	ongoing
3. Monitor additional 16 sites in the lakes (Arrowhead-4, Camelot-8, Sherwood-4) in the lakes and 2 toes drains at each dam.	Adams LWCD Tri-Lakes Mgmt District Lake Associations	ongoing
4. Work with the WDNR to learn new solutions as they become available. Utilize WDNR website as well as Lake Specialist at Adams LWCD.	Tri-Lakes Mgmt District	ongoing
5. Evaluate water quality report produced by UWSP & identify action plan.	Tri-Lakes Mgmt District	2015-2016
Educate community and public about ways to improve water quality.		
1. Write and publish informational articles in lake association newsletters, and .in Wis Rapids Tribune. Also post to websites & distribute mterials at public events & Tri-Lakes Annual Meeting. Combine family activity days to reach a wider & younger audiences.	Adams LWCD Lake Associations Tri-Lakes Mgmt District	ongoing
2. Distribute landowner packet of information prepared by Adams County LWCD to waterfront property owners.	Tri-Lakes Mgmt District	ongoing
3, Educate public about proper disposal of prescription drugs.	Tri-Lakes Mgmt District	ongoing
4. Establish website to serve as portal of information to the proper owners.	Tri-Lakes Mgmt District	2014 completed
Practice proper land use utilizing Comprehensive Plans and Zoning		
1. Within the Tri-Lakes watershed, continue to implement existing township comprehensive plan. Planning Commmission will be reviewing regularly.	Town of Rome Tri-Lakes Mgmt District	annually
2. Encourage smart growth plan development by attending town meetings in those towns within the Tri-Lakes watershed who do not have comprehensive plans.	Adams County P & Z	ongoing

<u>LAKE LEVELS: GOALS and ACTION ITEMS</u>	<u>WHO</u>	<u>WHEN</u>
Maintain lake levels that enhance water quality and meet the requirements of Wisconsin Statute Chapter 31.		
1. Operate dams to maintain lake levels and outflows as stated by WDNR. These levels are These operating levels are listed in Adams County I.O.M Plan (Inspection, Operation & Maintenance Plan). See Appendix G	Adams LWCD	ongoing
2. Investigate groundwater net flow around and through dams and determine what nutrient content the flow contains. Take water samples from toe drains for phosphorus content. Present findings to WDNR.	Tri-Lakes Mgmt Dist. Adams LWCD State Hygiene Lab	ongoing
Maintain stable stream flows into and out of the Tri-Lakes.		
1. Operate dams in proactive manner so large quantities of water are not released causing downstream flooding & streambank erosion. Lake levels will be lowered in a slow manner to accommodate anticipated heavy flows from the upper watershed.	Adams LWCD	annually
2. Develop plan to monitor flows.	Tri-Lakes Mgmt District Adams LWCD UWSP, WDNR	ongoing
3 Work with Central Sands water resource group.	Tri-Lakes Mgmt District	ongoing
<u>WATERSHED: GOALS and ACTION ITEMS</u>	<u>WHO</u>	<u>WHEN</u>
Reduce 50% the total phosphorus, total nitrogen, and total solid sediments entering Leola Ditch, 14 Mile Creek, and Unnamed Ditch 13-13 by 2020.		
1. Develop strategy to implement buffer corridor compliance. Maintain & monitor the buffer corridors.	Adams LWCD Adams P & Z	completed
2. Invite the Leola Drainage District governing body & Adams County Drainage Board to attend a joint meeting and/or request tour of the ditches..	Tri-Lakes Mgmt District	ongong
3. Investigate and discuss methods to purchase land in the watershed in order to broker land use change.	Tri-Lakes Mgmt District	2016-2017

4. Sample & test water at the headwaters region of the drainage district & as it enter Lake Camelot to determine clarity, dissolved oxygen, pH, temperature, biological conditions, total phosphorus, total nitrogen and total sediments.	Tri-Lakes Mgmt District Adams LWCD	ongoing
5. Implement the Adams County Stormwater Runoff ordinance to prevent and/or reduce nutrients & sediments from entering the state's waters.	Adams LWCD	ongoing
6. Monitor & advocate change in Drainage District legislation.	Tri-Lakes Mgmt District	ongoing
7. Implement State Agricultural Performance Standards by contacting agricultural producers to offer cost share for compliance with preventing runoff from livestock confinement operation and uncontained livestock manure; sedimentation due to soil erosion from croplands and stream banks; livestock access to stream sites.	Tri-Lakes Mgmt District Adams LWCD	ongoing
Create community awareness and improve community knowledge about watershed activities		
1. Educate landowners, agricultural producers, and community about State Agricultural Performance Standards and best management practices utilizing newsletters, meetings, and individual visits.	Adams LWCD, NRCS DNR, Waushara LWCD CREP	ongoing