Jeanne, scherere wisconsir gov Data Form 608-275-3283

AIS Early Detection Monitoring Data Form

Form 3200-xxx (R 6/2013)

Lake Name	County	WBIC	AIS sign?	Secchi (ft or m)	Conductivity (ZM tow if > 99 umhos/cm)
Hemilar	Sunc	1286100	(A) N	4	
Date(s)	Data collectors	Start time (nearest 15	min)	End time (nearest 15 min)	Total collector time (hrs x # collectors)
710/3	Jeanne Scherer	12:45		3:00 2:45	thr

Look for the following species: Purple loosestrife, Phragmites, flowering rush, Hydrilla, Brazilian waterweed, Eurasian water-milfoil, curly-leaf pondweed, yellow floating heart, zebra mussel, quagga mussel, Chinese mystery snail, banded mystery snail, faucet snail, New Zealand mud snail, didymo, water flea, and any other AIS found.

STEP 1: Record locations of sampling sites (in decimal degrees). Sampling sites include all public boat landings (BL), 5 targeted sites (TS) and the meander survey sites (MS). List AIS found at each site or record none. Collect a sample of any new AIS found. Collect five new invasive plant specimens, 20 Dreissenids, and 30 of each snail species and label with species, collector, date, lake name, WBIC and sampling site.

	Site	Latitude	Longitude	Snorkel (Y or N*)	If N snorkel, indicate why	Species, density 1-5 <sup>‡</sup>	
V	BL	43.59/04	90,14297	N	to Jalgar	CME (3) (EARL)	
Vms	建	43,59042	90 149300		and the second	CLP(3)	
V	151	43 58944	90.14414	N	heavy wortall	CLP(3)	
. 1	mg2	43.58952	90.14475		J	CLP (2)	
1	152	43.5898D	90 14520	N		00	
	153	4359009	90,4699	N		CLP 1	-
/	M 53	4358881	90 14 600	not p	utinswing	Photo Ferrestrial plant	
V	:m4	4358867	90.14526		4	latential yellow ms	nobloom
-	ms9	435 8847	9.0 14466	. 2		· · · · · · · · · · · · · · · · · · ·	
-	134	43.58967	90,14195	R	turbid	CLP(2)	
les and	135	43,591011	90.14209			Potential y. Man iris (1), CLR3) (	ms(1)
				2			7 Gath
		,			ŝ	1-1-1-1	12
				-			-

## \*For lakes/sites not snorkeled, substitute:

Boat landing site - 15 rake throws and 15 D-net samples OR 30 minutes, whichever comes first

Targeted site - 5 rake throws and 5 D-net samples OR 10 minutes, whichever comes first

50 meander sites - 10 rake throws and 10 D-net samples during meander survey between sampling sites for a total of 50 meander survey sites

†If lake/site was not snorkeled, indicate why: stained water, turbid water, blue-green bloom, chemical treatment other (please describe).



- 1 A few plants or invertebrates
- 2 One or a few plant beds or colonies of invertebrates
- 3 Many small beds or scattered plants or colonies of invertebrates

- 4 Dense plant, snail or mussel growth in a whole bay or portion of the lake
- 5 Dense plant, snail or mussel growth covering most shallow areas

Step 2: Collect Waterflea Tows from 3 sites: the deep hole (DH) and 2 other sites in water deeper than 15 feet (if possible). Submit sample and Water Flea To Monitoring Reprt form to Science Services.

Site	Depth sampled	Method (hor, obliq, vert)	Net diameter (30 or 50 cm)	Ethanoi added (Y or N)	Samples combined (Y or N)	Sample sent to, date
DH'		Kovizontal	CZ, O	У	Y	7/25/13
DHI	1	(~ ;	<b>3</b> 0	7	. 4	10
DHZ	45		<b>E</b> D	Y	'V	8 6

Step 3: Collect Veliger Tows from 3 sites; the deep hole (DH), outlet site (OS), and or downwind site (DS) in water depth of about 4 meters (if possible). Submit sample and Mussel Veliger Tow Monitoring Report form to Science Service.

Site	Depth sampled	Net diameter (30 or 50 cm)	Ethanol added (Y or N)	Samples combined (Y or N)	Sample sent to, date
DH(	5	<b>5</b> 0	~\\\\	У	7/25/12
DH1	5.	<b>5</b> 0	X	¥	41
DHZ	5	\$,0	NO	7	£ 23x

Notes: