			1 00000		
Data Collectors Sam Betterley, Erin Vinnie-Vollrath, Scott Egeren Date 7/5/12					
Lake Name Lost Lake		County	WBIC 140700	00	
Start Time 14:00 End Time	Secchi Dep	th feet or meters (circle	cone) Conductivity US/cm		
Look for the following species: Purple loosestrife, Pl					
heart, zebra mussel, quagga mussel, Chinese myster 50 rake and D-net samples during meander survey.				tes not snorkeled, take	
Did you snorkel the search sites? Y/N	If not, why? (circle o	one) stained water, turbid w	ater, blue-green bloom, chemical treati	ment, ot <mark>he</mark> r	
Rake/D-net counts: Species 1	Count; S	pecies 2 Cour	nt; Species 3 Cou	unt;	
Species 4	Count; S	pecies 5 Coun	t; Species 6 Co	unt	
STEP 1: Record locations of sites (in decimal de suspected AIS found.	grees) using a GPS (unit (datum WGS84). List AIS	found at each site or record none. Colle	ect a sample of any	
Boat Landing# Species CMS	<u> </u>	atitude 44.84883	Longitude89,33897	Density (1-5)	
Boat Landing# Species		atitude	Longitude	Density (1-5)	
Search Site# Species	<u> </u>	atitude <u>44,84970</u>	Longitude 89,3385/	Density (1-5)	
Search Site# 2 Species	, <u> </u>	atitude 44,85106	_Longitude89,33978	Density (1-5)	
Search Site# 3 Species	1	atitude 44,85 so 4	_Longitude89,34527	Density (1-5)	
Search Site# 4 Species CMS		atitude 44.84905	Longitude -89.34301	Density (1-5)	
Search Site# 5 Species	i	_atitude44,8467	Longitude	Density (1-5)	
Search Site# Species	L	_atitude	Longitude	Density (1-5)	
Meander Survey# Species	, L	_atitude	Longitude	Density (1-5)	
Meander Survey# Species		_atitude	Longitude	Density (1-5)	
Meander Survey# Species	2 [_atitude	Longitude	Density (1-5)	

Step 2: Label first five specimens collected with species, collector, date, lake name, WBIC and Location # Send your specimens to an expert for verification. Instructions on how to voucher specimens and a list of statewide taxonomy experts can be found at: http://dnr.wi.gov/invasives/aquatic/whattodo/staff/

	Step 3: Collect Waterliea Tows from three sites around the lake in water deeper than 15 feet (if possible).
	Method used: horizontal tows (near surface) or oblique tows (near bottom to surface if greater than 15 feet)
	Diameter of plankton net mouth (circle one) 30cm (50cm)other
	Depth sampled: Tow 1 15 ft Tow 2 15 ft Tow 3 15 ft
William I	Depth sampled: Tow 1 15 ft Tow 2 15 ft Tow 3 15 ft Has ethanol been added? Y/N Have samples been consolidated into one bottle? Y/N
	a built for law
	Step 4: Collect Veliger Tows from three sites in 5-10 feet of water (within a meter of the bottom) Conductivity too low
·	Guidelines: If Secchi depth is >4m take two 2m deep samples; if Secchi is between 2-4m take one 2m deep sample; if Secchi is <2m take one 1m tow.
	Diameter of plankton net mouth (circle one) 30cm 50cm other
	Has ethanol been added? Y/N Have samples been consolidated into one bottle? Y/N
	Step 5: Data was entered into SWIMS on 10-16-2017 by Sam Betterley
	Date V Name
	, , , , , , , , , , , , , , , , , , ,

Density Ratings

- 1 A few plants or invertebrates
- 2 One or a few plant beds or colonies of invertebrates
- 3 Many small beds or scattered plants or colonies of invertebrates
- 4 Dense plant, snail or mussel growth in a whole bay or portion of the lake
- 5 Dense plant, snail or mussel growth covering most shallow areas

General guidance on areas to search for the 10 minute guick snorkel search sites:

- Check rocks for zebra/quagga mussels, faucet snails and New Zealand mudsnails.
- · Check around small backyard boat launches.
- Check near creek inlets (especially if AIS are found upstream).
- Check the stems of emergent vegetation for climbing faucet snails.
- Check areas downwind of large boat landings.



Present-CMS ZM - not suitable