
Presque Isle Wilderness Waters Program

Aquatic Plant Management Plan – Crab Lake

Prepared for:

Presque Isle Town Lakes Committee
Contact: Otto Novak
P.O. Box 37
Presque Isle, Wisconsin 54557

Prepared by:

White Water Associates, Inc.
Dean Premo, Ph.D.
429 River Lane, P.O. Box 27
Amasa, Michigan 49903



Date: December 2015

Page left intentionally blank



Presque Isle Wilderness Waters Program

Aquatic Plant Management Plan – Crab Lake

This plan is a product of WDNR Lake Planning Grant (LPL-1508-13) awarded to:

Presque Isle Town Lakes Committee
P.O. Box 37
Presque Isle, Wisconsin 54557
Contact: Otto Novak
Phone: (715) 686-2628; E-mail: wefish2@hughes.net

Submitted to:

Wisconsin Department of Natural Resources
Attention: Kevin Gauthier, Sr., Water Resource Management Specialist
8770 Hwy J
Woodruff, WI 54568
Phone: (715) 365-5211 ext. 214; Email: Kevin.GauthierSr@wisconsin.gov

Prepared by:

White Water Associates, Inc.
Dean Premo, Ph.D., Angie Stine, B.S., Caitlin Clarke, B.S., and Kent Premo, M.S.
429 River Lane, P.O. Box 27
Amasa, Michigan 49903
Phone: (906) 822-7889; E-mail: dean.premo@white-water-associates.com

Cite as: Premo, Dean, Angie Stine, Caitlin Clarke and Kent Premo. 2015. Presque Isle Wilderness Waters Program: Crab Lake Aquatic Plant Management Plan. White Water Associates, Inc.



Page left intentionally blank



TABLE OF CONTENTS

Chapter 1. Introduction	1
Chapter 2. Study Area	4
Chapter 3. Purpose and Goal Statements	8
Chapter 4. Information and Analysis	9
Part 1. Watershed	9
Part 2. Aquatic Plant Management History	13
Part 3. Aquatic Plant Community Description	13
Part 4. Fish Community	17
Part 5. Water Quality and Trophic Status	18
Part 6. Water Use	18
Part 7. Riparian Area	18
Part 8. Wildlife	19
Part 9. Stakeholders	20
Chapter 5. Recommendations, Actions, and Objectives	21
Chapter 6. Contingency Plan for AIS	25
Appendix A – Literature Cited	
Appendix B – Tables and Figures	
Appendix C – Review of Lake Water Quality	

Page left intentionally blank



CHAPTER 1

Introduction

The *Presque Isle Wilderness Waters Program* results from the efforts of the Presque Isle Town Lakes Committee, an organization that has been active since 2005. The Lakes Committee views stewardship of lakes as an ongoing endeavor that is integrated, coordinated, and administered by the Lakes Committee. This broader perspective accommodates the appropriate range of geographic scales from which to approach lake stewardship: a discrete “lake specific” focus that goes hand-in-hand with waterscape-wide awareness.

This aquatic plant management plan addresses Crab Lake. Despite this specificity, it maintains the waterscape perspective crucial to effective lake stewardship. This is especially important when it comes to preventing introduction and establishment of aquatic invasive species (AIS). The closely related *Wilderness Waters Adaptive Management Plan* (Premo et al., 2015) provides additional overarching waterscape level examination that allows greater opportunity and efficiency in water resource management and education.

A systematic survey of aquatic plants using the Wisconsin Department of Natural Resources (WDNR) “point-intercept” method was an important underpinning of this aquatic plant management plan. An analysis of the plant data along with water quality and other lake information allowed the preparation of the plan.

Aquatic plants rarely get the respect they merit, although this is slowly changing. We still call an aquatic plant bed a “weed bed.” Many aquatic plants have “weed” in their names (e.g., duckweed, pondweed, or musky weed). Likely this term was borrowed from “seaweed” and not intended as derogatory, but in today’s use, “weed” connotes an unwanted, aggressively growing plant. Such is not the case for the vast majority of aquatic plants. In fact, aquatic plants are a vital part of a lake ecosystem, recycling nutrients, providing vertical and horizontal structure, and creating habitat for animal life. Invertebrates, including crustaceans and insects, live on or within this “aquatic forest.” Fish find food and shelter within aquatic plant beds. Waterfowl eat parts of plants directly as well as feed on invertebrates associated with the plants. Muskrats eat aquatic plants and particularly love cattails and bulrushes. Otter and mink hunt invertebrates and small vertebrates within the shelter of submergent and emergent beds. In shallow water, great blue herons find fishes among the plants.

In lakes that receive an excess of nutrients (particularly from fertilizers or leaking septic tanks), plant growth can become too lush or dominated by only a few species. As these abundant plants die, their decomposition can depress dissolved oxygen levels and diminish suitability for fish. Algae can respond rapidly to nutrient influxes and create nuisance conditions. These phenomena can cause humans to view all aquatic plants in a negative light.

On another negative front, non-native plant species, transported on boats and trailers or dumped from home aquariums, private ponds and water gardens may come to dominate a water body to the exclusion of a healthy diversity of native species. Eurasian water-milfoil (*Myriophyllum spicatum*) is one of the better known examples of these so-called aquatic invasive plant species.

For most lakes, native aquatic plants are an overwhelmingly positive attribute, greatly enhancing the aesthetics of the lake and providing good opportunities for fishing, boating, swimming, snorkeling, sight-seeing, and hunting.

When it comes to aquatic plant management, it is useful to heed the mantra of the medical profession: “First, do no harm.” It is both a social and scientific convention that aquatic plant management is more effective and beneficial when a lake is considered as an entire and integrated ecosystem. Anyone involved in aquatic plant management should be aware that a permit may be required to remove, add, or control aquatic plants. In addition, anyone using Wisconsin’s lakes must comply with the “Boat Launch Law” that addresses transport of aquatic plants on boat trailers and other equipment. A good review of the laws, permits, and regulations that affect management and behavior surrounding aquatic plants can be found in the WDNR guidelines called *Aquatic Plant Management in Wisconsin*.¹

In preparing this plan, we followed guidelines in *Aquatic Plant Management in Wisconsin*. The resulting plan is an adaptive plan (Walters, 1986). Simply put, it will be modified as new information becomes available. The WDNR Guidance document outlines three objectives that may influence preparation of an aquatic plant management plan. Currently, the motivation for this plan lies in the first two objectives:

- **Protection** - preventing the introduction of nuisance or invasive species into waters where these plants are not currently present;
- **Maintenance** - continuing the patterns of recreational use that have developed historically on and around a lake; and

¹ <http://www4.uwsp.edu/cnr/uwexlakes/ecology/APM/APMguideFull2010.pdf>

-
- **Rehabilitation** - controlling an imbalance in the aquatic plant community leading to the dominance of a few plant species, frequently associated with the introduction of invasive non-native species.

During projects with the WDNR Planning Grant Program and through past efforts, Town Lakes Committee has followed the first five steps in the seven-step plan outlined in the Guidance Document for developing an aquatic plant management plan:

1. Goal setting – Getting the effort organized, identifying problems to be addressed, and agreeing on the goals;
2. Inventory – Collecting baseline information to define the past and existing conditions;
3. Analysis – Synthesizing the information, quantifying and comparing the current conditions to desired conditions, researching opportunities and constraints, and setting directions to achieving the goals;
4. Alternatives – Listing possible management alternatives and evaluating their strengths, weaknesses and general feasibility;
5. Recommendations – Prioritizing and selecting preferred management options, setting objectives, drafting the plan;
6. Implementation – Formally adopting the plan, lining up funding, and scheduling activities for taking action to achieve the goals;
7. Monitor & Modify – Developing a mechanism for tracking activities and adjusting the plan as it evolves.

Besides this introductory chapter, this plan is organized in six Chapters. The study area is described in Chapter 2. Chapter 3 states the purpose and goals for the plan. Chapter 4 presents an inventory and analysis of information that pertain to the plan including the results of the aquatic plant survey. Chapter 5 provides recommendations that support the overall goals and establish the stewardship component of plan. Finally, Chapter 6 presents actions and objectives for implementing the plan. Three appendices complete this document. Appendix A contains literature cited, Appendix B contains tables and figures for the aquatic plant survey, and Appendix C contains a *Review of Crab Lake Water Quality*.


Page left intentionally blank



CHAPTER 2


Study Area

Presque Isle Township is one of the northern-most townships in Vilas County, Wisconsin. Presque Isle Township's northern border is shared with the State of Michigan. In fact some of the Presque Isle Township lakes lie on the state border. The location of the subject of this APM Plan (Crab Lake) is shown in Exhibit 1 along with other lakes in Presque Isle Township that have had point-intercept aquatic plant surveys conducted. Exhibit 2 is an aerial view of Crab Lake.



“Almost an island” is the literal translation of the French phrase “Presque Isle.” Early French missionaries, perhaps disoriented by the preponderance of water in this north central Wisconsin landscape applied the name, “Presque Isle” to describe an area where the water seemed to dominate the land. The French visitors and Native Americans certainly recognized this landscape as special. Modern ecologists and recreationist share this view. The region that includes the Township of Presque Isle, Wisconsin is an ecological landscape marvelously rich in surface waters. Aerial photography reveals a concentration of lakes and streams that is unique in North America. Presque Isle Township has eighty-four lakes. The Presque Isle area could as easily be termed a “waterscape” as a “landscape.”

Exhibit 2. Aerial photo of Crab Lake.




Descriptive parameters for Crab Lake are in Exhibit 3. It is a drainage lake of about 909 acres and maximum depth of 60 feet. The shoreline development index values for twenty-nine of the Wilderness Waters Program lakes surveyed from 2007 to 2014 ranged from 1.2 to 4.2

(average = 2.1). Crab Lake's shoreline development index is higher than this average (3.7). The shoreline development index is a quantitative expression derived from the shape of the lake. It is defined as the ratio of the shoreline length to the length of the circumference of a circle of the same area as the lake. A perfectly round lake would have an index of 1. Increasing irregularity of shoreline development in the form of bays and projections of the shore is shown by numbers greater than 1. For example, fjord lakes with extremely irregularly shaped shorelines sometimes have SDI's exceeding 5. A higher shoreline development index indicates that a lake has relatively more productive littoral zone habitat.

Exhibit 3. Water Body Parameters	
Water Body Name	Crab
County	Vilas
Township/Range/Section	T43N-R06E-S13,14,15,21,22,23,24
Water Body Identification Code	2953500
Lake Type	Drainage
Surface Area (acres)	909
Maximum Depth (feet)	60
Maximum Length (miles)	1.2
Maximum Width (miles)	1.5
Shoreline Length (miles)	15.8
Shoreline Development Index	3.7
Total Number of Piers (2013 aerial)	78
Number of Piers / Mile of Shoreline	4.9
Total Number of Homes (2013 aerial)	67
Number of Homes / Mile of Shoreline	4.3

Crab Lake has one public access site. We observe a total of 78 piers on the shoreline of Crab Lake from a 2013 aerial photograph or about 4.9 piers per mile of shoreline. The riparian

area consists of both upland and wetland areas (Exhibit 4). Crab Lake is considered an Outstanding Resource Water (ORW) (WI Admin. Code, 2010). An ORW is: a body of water that provides outstanding recreational opportunities, supports valuable fisheries and wildlife habitat, has good water quality, and is not significantly impacted by human activities. It does not typically have any point sources discharging pollutants directly into the water, though it may receive runoff from nonpoint sources. Also, new discharges may be permitted only if their effluent quality is equal to or better than the background water quality of that waterway at all times—no increases of pollutant levels are allowed (WDNR, 2013).



CHAPTER 3

Purpose and Goal Statements

This plan approaches aquatic plant management with a healthy dose of humility. We do not always understand the causes of environmental phenomena or the effects of our actions to manage the environment. With that thought in mind, we have crafted a statement of purpose and goals for this plan:

Crab Lake has a native aquatic plant community that was documented by a point-intercept aquatic plant survey. This plant community is essential to, and part of, a high quality aquatic ecosystem that benefits the human community with its recreational and aesthetic features. The purpose of this aquatic plant management plan is to maintain the aquatic plant community in its present high quality state.

Supporting this purpose, the goals of this aquatic plant management plan are:

- (1) Monitor and protect the native aquatic plant community;*
- (2) Prevent establishment of AIS and nuisance levels of native plants;*
- (3) Promote and interpret APM efforts; and*
- (4) Educate riparian owners and lake users on preventing AIS introduction, reducing nutrient inputs that potentially alter the plant community, and minimizing physical removal of native riparian and littoral zone plants.*

The purpose and goals are the foundation for the aquatic plant management plan presented in this document. They inform the objectives and actions outlined in Chapter 5 and are the principal motivation of Crab Lake stewards.

Page left intentionally blank




CHAPTER 4

Information and Analysis

Our efforts in the Wilderness Waters Program have compiled information about historical and current conditions of the Crab Lake ecosystem and its surrounding watershed. Of particular importance to this aquatic plant management plan is the aquatic plant survey that was conducted using the *WDNR Protocol for Aquatic Plant Survey, Collecting, Mapping, Preserving, and Data Entry* (Hauxwell et al., 2010). The results of this comprehensive “point-intercept” survey along with relevant components of other information are presented in this chapter under nine respective subheadings: watershed, aquatic plant management history, aquatic plant community description, fish community, water quality and trophic status, water use, riparian area, wildlife, and stakeholders.


Part 1. Watershed

The Presque Isle Township waterscape sits on a large-scale watershed divide. Some of the water drains north through the Presque Isle River system and eventually enters Lake Superior. Some of the water drains into the Wisconsin River system to the Mississippi River and to the Gulf of Mexico. In fact there are two federal hydrologic sub-basins (designated by 8-digit HUC codes) that include Presque Isle Township. The Black-Presque Isle Rivers sub-basin (HUC#04020101) drains north to Lake Superior and the Flambeau River sub-basin (HUC#0705002) drains southwesterly to the Mississippi River. The Black-Presque Isle Rivers sub-basin contains two federal hydrologic sub-watersheds within Presque Isle Township: the South Branch Presque Isle River sub-watershed (HUC#040201010303) and the Pomeroy Creek-East Branch Presque Isle River sub-watershed (HUC#040201010301). The Flambeau River sub-basin contains one sub-watershed within Presque Isle Township: the Rice Creek sub-watershed (HUC#07050020103). Exhibit 5 illustrates these watersheds and the watersheds of the water bodies subject to the Wilderness Waters Program studies. Crab Lake is contained within the South Branch Presque Isle River sub-watershed (Exhibit 5).



Description: Portions of 3 federal hydrologic sub-watersheds are illustrated: (1) S.Br. Presque Isle River (bounded by yellow), (2) Pomeroy Cr-E.Br. Presque Isle River (bounded by blue and yellow), and (3) Rice Cr. (bounded by green, yellow, and blue). Also shown are the smaller watersheds of individual water bodies subject to the Wilderness Waters studies (bounded by red and, in the case of Papoose Lake, bounded by white). White dots show outlet points for the smaller watersheds.

The elevation in Presque Isle Township ranges from around 1,550 feet above sea level to 1,750 feet above sea level. A digital elevation model is provided as Exhibit 6 and shows the relative elevations for the area with orange areas of the landscape being the highest elevations and greens and blues being the lowest elevations.



The watershed (drainage basin) is all of the land and water areas that drain toward a particular river or lake. A water body is greatly influenced by its watershed. Watershed size, topography, geology, land use, soil fertility and erodibility, and vegetation are all factors that influence water quality. The Crab Lake watershed is about 12,990 acres. The type of land cover (for example, forest, grassland, row crops, or human development) is also an important variable in determining amounts and kinds of materials (like nutrients and sediment) that are carried off the land and into the water. Certain kinds of agriculture (tilled row crops) and urban areas (with their impervious surfaces) have a tendency to give up sediments and nutrients to runoff. In contrast, native vegetation (forests, wetlands, and grasslands), tend to slow runoff of water and nutrients, allowing the soil to absorb them. The cover types in Crab Lake's watershed are presented in Exhibit 7.

Exhibit 7. Cover Types and Soil Groups of the Crab Lake Watershed.			
Cover Type		Acres	Percent
Agriculture		0	0
Commercial		0	0
Forest		7163.3	55.1
Grass/Pasture		9.1	0.1
High-density Residential		0	0
Low-density Residential		845.8	6.5
Water		4972.7	38.3
Total		12991.1	100.0
Soil Group	Acres	Percent	Hydrologic Soil Groups - Soils are classified by the Natural Resource Conservation Service into four Hydrologic Soil Groups* based on the soil's runoff potential. The four Hydrologic Soils Groups are A, B, C and D. Where A has the smallest runoff potential and D the greatest.
A	956.0	7.4	Group A is sand, loamy sand or sandy loam types of soils. It has low runoff potential and high infiltration rates even when thoroughly wetted. They consist chiefly of deep, well to excessively drained sands or gravels and have a high rate of water transmission.
B	6957.5	53.6	Group B is silt loam or loam. It has a moderate infiltration rate when thoroughly wetted and consists chiefly or moderately deep to deep, moderately well to well drained soils with moderately fine to moderately coarse textures.
C	361.8	2.8	Group C soils are sandy clay loam. They have low infiltration rates when thoroughly wetted and consist chiefly of soils with a layer that impedes downward movement of water and soils with moderately fine to fine structure.
D	4715.8	36.3	Group D soils are clay loam, silty clay loam, sandy clay, silty clay or clay. This soil has the highest runoff potential. They have very low infiltration rates when thoroughly wetted and consist chiefly of clay soils with high swelling potential, soils with a permanent high water table, soils with a claypan or clay layer at or near the surface and shallow soils over nearly impervious material.

*(USDA, Natural Resources Conservation Service, 1986)

Forest and surface water comprise the largest components. Soil group B is most prevalent, followed by group D, while groups A and C make up a small percentage of the watershed. Soil group A has the highest infiltration capacity, and the lowest runoff potential. Conversely, soil group D has the lowest infiltration capacity, and the highest runoff potential. The watershed to

lake area ratio is 14:1. Water quality often decreases with an increasing ratio of watershed area to lake area. As the watershed to lake area increases there are more sources and amounts of runoff. In larger watersheds, runoff water can leach more minerals and nutrients and carry them to the lake. The runoff to a lake (such as after a rainstorm or snowmelt) differs greatly among land uses. Forest cover is the most protective as it exports much less soil (through erosion) and nutrients (such as phosphorus and nitrogen) to the lake than agricultural or urban land use.

Part 2. Aquatic Plant Management History

As far as we can determine, no systematic or large-scale plant management activity has ever taken place in Crab Lake. Over the years, no particular nuisance issues have demanded control action. In 2007, an aquatic plant survey was conducted on Crab Lake, and again in the summer of 2014. Findings from the 2007 and 2014 surveys are discussed in the next section (Part 3).

Part 3. Aquatic Plant Community Description

Why do lakes need aquatic plants? In many ways, they are underwater forests. Aquatic plants provide vertical and horizontal structure in the lake just like the many forms and variety of trees do in a forest. Imagine how diminished a forest's biodiversity becomes in the advent of a clear-cut. Similarly, a lake's biodiversity in large part depends on a diversity of plants.

Aquatic plants are beneficial in many ways. Areas with plants produce more food for fish (insect larvae, snails, and other invertebrates). Aquatic vegetation offers fish shelter and spawning habitat. Many submerged plants provide food for waterfowl and habitat for insects on which some waterfowl feed. Aquatic plants further benefit lakes by producing oxygen and absorbing nutrients (phosphorus and nitrogen) from runoff. Aquatic plants also protect shorelines and lake bottoms by dampening wave action and stabilizing sediments.

The distribution of plants within a lake is generally limited by light availability, which is, in turn, controlled by water clarity. Aquatic biologists often estimate the depth to which rooted aquatic plants can exist as about two times the average Secchi clarity depth. For example, if the average Secchi depth is eight feet then it is fairly accurate to estimate that rooted plants might exist in water as deep as sixteen feet. At depths greater than that (in our hypothetical example), light is insufficient for rooted plants to grow. In addition to available light, the type of substrate influences the distribution of rooted aquatic plants. Plants are more likely to be found in muddy or soft sediments containing organic matter, and less likely to occur where the substrate is sand,

gravel, or rock. Finally, water chemistry influences which plants are found in a body of water. Some species prefer alkaline lakes and some prefer more acidic lakes. The presence of nutrients like phosphorous and nitrogen also influence plant community composition.

As mentioned earlier, non-native invasive plant species can reach high densities and wide distribution within a lake. This diminishes the native plant community and the related habitat. At times, even a native plant species can reach nuisance levels with respect to certain kinds of human recreation. These cases may warrant some kind of plant management.

Aquatic plant surveys have been conducted on Crab Lake by aquatic plant specialists in 2007 and 2014. In each of these surveys, WDNR point-intercept protocol and methodology was followed. This formal survey assesses the plant species composition on a grid of several hundred points distributed evenly over the lake. Using latitude-longitude coordinates and a handheld GPS unit, scientists navigate to the points and use a rake mounted on a pole or rope to sample plants. Plants are identified, recorded and put into a dedicated spreadsheet for storage and data analysis. This systematic survey provides baseline data about the lake. Available data from the 2007 survey includes species and relative frequency. A comparison of these data is presented below. Supporting tables and figures for the 2014 aquatic plant survey is provided in Appendix B. Table 3 provides a comparison of statistics from both survey years.

Species richness refers to the total number of species recorded. We recorded 35 species of aquatic plants. Of these, 29 were collected at sampling sites and the others were observed from the boat. Table 1 displays summary statistics for the survey. Table 2 provides a list of the species encountered, including common and scientific name along with summarizing statistics.² The number of species encountered at any given sample point ranged from 0 to 7 and 58 sample points were found to have aquatic vegetation present. The average number of species encountered at these vegetated sites was 1.93. The actual number of species encountered at each of the vegetated sites is graphically displayed on Figure 1. Plant density is estimated by a “rake fullness” metric (3 being the highest possible density). These densities (considering all species) are displayed for each sampling site on Figure 2.

The maximum depth of plant colonization is 10.5 feet (Table 1 and Figure 3). Rooted vegetation was found at 58 of the 239 sample sites with depth \leq the maximum depth of plant colonization (24.27% of sites). These sites are displayed as a black dot within a circle on Figure 4. This indicates that although availability of appropriate depth may limit the distribution of

² If you are interested in learning more about the plant species found in the lake, visit the University of Wisconsin Steven Point Freckmann Herbarium website at: <http://wisplants.uwsp.edu/> or obtain a copy of “Through the Looking Glass (A Field Guide to the Aquatic Plants in Wisconsin).”

plants, it is not the only habitat factor involved. Substrate is another feature that influences plant distribution (e.g., soft substrate often harbors more plants than hard substrate). Figure 5 presents the substrates encountered during the aquatic plant survey (mud, sand, or rock).

Table 2 provides information about the frequency of occurrence of the plant species recorded in the lake. Several metrics are provided, including total number of sites in which each species was found and frequency of occurrence at sites \leq the maximum depth of rooted vegetation. This frequency metric is standardized as a “relative frequency” (also shown in Table 2) by dividing the frequency of occurrence for a given species by the sum of frequency of occurrence for all plants and multiplying by 100 to form a percentage. The resulting relative frequencies for all species total 100%. The relative frequencies for the plant species collected with a rake are graphically displayed in descending order on Figure 6. This display shows that fern pondweed (*Potamogeton robbinsii*) had the highest relative frequency followed by wild celery (*Vallisneria americana*). The lowest relative frequencies are at the far right of the graph. As examples of individual species distributions, we show the occurrences of a few of the most frequently and least frequently encountered plants in Figures 8-13.

Species richness (total number of plants recorded at the lake) is a measure of species diversity, but it doesn’t tell the whole story. As an example, consider the plant communities of two hypothetical ponds each with 1,000 individual plants representing ten plant species (in other words, richness is 10). In the first pond each of the ten species populations is comprised of 100 individuals. In the second pond, Species #1 has a population of 991 individuals and each of the other nine species is represented by one individual plant. Intuitively, we would say that first pond is more diverse because there is more “even” distribution of individual species. The “Simpson Diversity Index” takes into account both richness and evenness in estimating diversity. It is based on a plant’s relative frequency in a lake. The closer the Simpson Diversity Index is to 1, the more diverse the plant community. The Simpson Diversity Index for Crab Lake aquatic plants is 0.92 (Table 1) which indicates a highly diverse aquatic plant community.

Another measure of floristic diversity and quality is the *Floristic Quality Index* (FQI). Floristic quality is an assessment metric designed to evaluate the closeness that the flora of an area is to that of undisturbed conditions (Nichols, 1999). Among other applications, it forms a standardized metric that can be used to compare the quality of different lakes (or different locations within a single lake) and monitor long-term changes in a lake’s plant community (an indicator of lake health). The FQI for a lake is determined by using the average *coefficient of conservatism* times the square root of the number of native plant species present in the lake.

Knowledgeable botanists have assigned to each native aquatic plant a *coefficient of conservatism* representing the probability that a plant is likely to occur in pristine environments (relatively unaltered from presettlement conditions). The coefficients range from 0 to 10, with 10 being assigned to those species most sensitive to disturbance. As more environmental disturbance occurs, the less conservative species become more prevalent.


Nichols (1999) analyzed aquatic plant community data from 554 Wisconsin Lakes to ascertain geographic (ecoregional) characteristics of the FQI metric. This is useful for considering how the Crab Lake FQI (37.2) compares to other lakes and regions. The statewide medians for number of species and FQI are 13 and 22.2, respectively. Crab Lake values are significantly higher than these statewide values. With regard to the Presque Isle Township lakes, the minimum FQI value is 17.1 (Armour Lake), the maximum is 39.8 (Morton Lake), and the average FQI value is 29.5. Compared to other lakes in Presque Isle Township, Crab Lake's FQI is much higher than the average (37.2). Nichols (1999) determined that there are four ecoregional-lake types groups in Wisconsin: (1) Northern Lakes and Forests lakes, (2) Northern Lakes and Forests flowages, (3) North Central Hardwoods and Southeastern Till Plain lakes and flowages, and (4) Driftless Area and Mississippi River Backwater lakes. Crab Lake is located in the Northern Lakes and Forests lakes group. Nichols (1999) found species numbers for the Northern Lakes and Forests lakes group had a median value of 13. Crab Lake data is much higher than that find. Finally, the Crab Lake FQI (37.2) is significantly higher than the median value for the Northern Lakes and Forests lakes group (24.3). This analysis supports the contention that the Crab Lake plant community is healthy and diverse and indicative of an undisturbed environment.

We found no state or federally listed species. We observed no aquatic plants in Crab Lake that would be considered a nuisance-level population density/distribution. We did observe a small patch of *Phragmites* (this had been previously reported on the WDNR website). We collected a specimen and delivered it to the UWSP herbarium. Dr. Freckmann said it appeared to be the native *Phragmites*. This patch deserves some further monitoring and sampling to verify this observation. We also observed some stands of cattails that should be further monitored and species determined.

Exhibit 8 displays the relative frequency of plants found in 2007 and 2014. It should be noted that in the 2007 survey, filamentous algae and aquatic moss were included in data collection, and each had a calculated relative frequency. The current WDNR protocol does not including these species, so the 2014 data did not calculate their relative frequencies. Because of

this difference in the 2007 and 2014 calculation of plant statistics, the relative frequencies of observed plants are not directly comparable.

We can see that many of the previously dominant species have decreased in frequency. Fern pondweed, on the other hand, has increased more than double. Also noted are the absence of many species from 2007 to 2014, and a large increase of new species not seen in 2007.



Part 4. Fish Community

It was beyond the scope of the current Wilderness Waters project to characterize the fish community and fish habitat of this water body. The WDNR Lake Pages website (<http://dnr.wi.gov/lakes/lakepages/>) indicates that the bottom is comprised of 50% sand, 30% gravel, 20% rock, and 0% muck and that fish species present include musky, panfish, largemouth bass, smallmouth bass, northern pike, and walleye.

Part 5. Water Quality and Trophic Status

Crab Lake is a 909 acre drainage lake with a maximum depth of 60 feet. Existing water quality information includes data was retrieved from the Wisconsin DNR SWIMS database between 1992 and 2014. On November 13, 2012 UW Stevens Point monitored various water quality parameters on Crab Lake. Secchi disk measurements were collected by Citizen Lake Monitoring Network (CLMN) volunteers from 1992 to 2014. The water quality information is briefly summarized in this section, but more fully interpreted in Appendix C.

Temperature and dissolved oxygen showed stratification in Crab Lake in the ice-free season. Average water clarity is considered “fair.” User perception of Crab Lake aesthetic quality is generally regarded as “beautiful, could not be nicer.” The trophic state is mesotrophic. Water quality would be classified as “very good” with respect to phosphorus concentrations. Chlorophyll *a*, turbidity, nitrogen, chloride, alkalinity, calcium, magnesium, sodium, and potassium are considered low. Crab Lake is also considered an Outstanding Resource Water (WI Admin. Code, 2010).

Part 6. Water Use

Crab Lake has one public access site, and is used by riparian owners and recreationists for a variety of activities. A small part of the southwestern shoreline is owned by the State of Wisconsin, along with four state-owned islands.

Part 7. Riparian Area

Part 1 (Watershed) describes the larger riparian area context of Crab Lake. The near shore riparian area can be appreciated by viewing Exhibits 2 and 4. These images give the impression that the lake is lightly developed with a fairly intact forested riparian zone that extends for hundreds of feet back from the lake. The forest is a mixture of coniferous and deciduous trees and shrubs. Our review of 2013 aerial photography reveals 67 houses on the lake. This intact riparian area provides numerous important functions and values to the lake. It effectively filters runoff to the lake. It provides excellent habitat for birds and mammals. Trees that fall into the lake from the riparian zone contribute important habitat elements to the lake. Educating riparian owners as to the value of riparian areas is important to the maintenance of these critical areas.

Part 8. Wildlife

A study of wildlife was beyond the scope of the current study, but would be valuable to study and interpret in future iterations of the plan. This would be especially true of wetland and water oriented wildlife such as frogs, waterfowl, fish-eating birds, aquatic and semi-aquatic mammals, and invertebrate animals. In the future, it would be desirable to monitor indicator species of wildlife such as common loons, bald eagles, and osprey. Also of special importance would be monitoring the populations of aquatic invasive animal species that already exist in the lake (rusty crayfish) and monitoring for the presence of currently not present aquatic invasive wildlife species (for example, spiny water flea, or zebra mussel) and fish species (for example, rainbow smelt or common carp).

Crab Lake is currently designated as a *priority navigable water* (PNW) and an *area of special natural resource interest* (ASNRI) (WI Admin. Code, 2014). Priority Navigable Waters meet any of these standards: navigable waterways, or portions thereof, that are considered ORW/ERW or trout streams; lakes less than 50 acres in size; tributaries and rivers connecting to inland lakes containing naturally-reproducing lake sturgeon populations; waters with self-sustaining walleye populations in ceded territories; waters with self-sustaining musky populations; or perennial tributaries to trout streams (WI Admin. Code, 2014). Crab Lake is considered a PNW because it is an ORW and has self-sustaining musky and walleye populations.

A water body designated as an Area of Special Natural Resource Interest can be any of the following: WDNR trout streams; Outstanding or Exceptional Resource Waters (ORW/ERW); waters or portions of waters inhabited by endangered, threatened, special concern species or unique ecological communities; wild rice waters; waters in ecologically significant coastal wetlands along Lake Michigan and Superior; or federal or state waters designated as wild or scenic rivers (WI Admin. Code, 2014). Crab Lake is considered an ASNRI because it is an Outstanding Resource Water and because it inhabits state or federally designated threatened or endangered species. The Wisconsin Natural Heritage Inventory (NHI) lists plants and animals as rare or sensitive species and/or communities that are considered high-quality and significant natural features (Exhibit 9). The following rare or sensitive species or communities are found in the same town/range as Crab Lake (NHI, 2014).

Exhibit 9. Rare Species and Communities located near Crab Lake.

<i>Common Name</i>	<i>Scientific Name</i>	<i>State Status</i> ³	<i>Group Name</i>
A predaceous diving beetle	<i>Agabus wasastjerna</i>	SC/N	Beetle
Trumpeter swan	<i>Cygnus buccinators</i>	SC/M	Bird
Boreal rich fen		NA	Community
Emergent marsh-wild rice		NA	Community
Ephemeral pond		NA	Community
Lake-deep, soft, seepage		NA	Community
Lake-spring		NA	Community
Northern mesic forest		NA	Community
Northern wet forest		NA	Community
Northern wet-mesic forest		NA	Community
Poor fen		NA	Community
Downy willow-herb	<i>Epilobium strictum</i>	SC	Plant
Fairy slipper	<i>Calypso bulbosa</i>	THR	Plant
Smith's melic grass	<i>Melica smithii</i>	END	Plant

Part 9. Stakeholders

At this juncture in the ongoing aquatic plant management planning process, the Town Lakes Committee has represented the Crab Lake stakeholders. Additional stakeholders and interested citizens are invited to participate as the plan is refined and updated in order to broaden input, build consensus, and encourage participation in stewardship. No contentious direct plant management actions (for example, harvesting or use of herbicides) are a component of the current plan.

³ **END**=Endangered; **THR**=Threatened; **SC**=Special Concern; **SC/P**=fully protected; **SC/N**=no laws regulating use, possession or harvesting; **SC/H**=take regulated by establishment of open/closed seasons; **SC/FL**=federally protected as endangered or threatened, but not so designated by DNR; **SC/M**=fully protected by federal and state laws under Migratory Bird Act.

CHAPTER 5

Recommendations, Actions, and Objectives

In this chapter we provide recommendations for specific objectives and associated actions to support the APM Plan’s goals stated in Chapter 3 and re-stated here for convenient reference:

- (1) Monitor and protect the native aquatic plant community;*
- (2) Prevent establishment of AIS and nuisance levels of native plants;*
- (3) Promote and interpret APM efforts; and*
- (4) Educate riparian owners and lake users on preventing AIS introduction, reducing nutrient inputs that potentially alter the plant community, and minimizing physical removal of native riparian and littoral zone plants.*

Since Crab Lake is a healthy ecosystem, we could simply recommend an alternative of “no action.” In other words, Crab Lake continues without any effort or intervention on part of lake stewards. Nevertheless, we consider the “no action” alternative imprudent. Many forces threaten the quality of the lake and Wilderness Waters Program and Town Lakes Committee feels a great responsibility to minimize the threats. We therefore outline in this section a set of actions and related management objectives that will actively engage lake stewards in the process of management.

The actions are presented in tabular form. Each “action” consists of a set of four statements: (1) a declarative “action” statement that specifies the action (2) a statement of the “objective” that the action serves, (3) a “monitoring” statement that specifies the party responsible for carrying out the action and maintaining data, and (4) a “status” statement that suggests a timeline/calendar and indicates status (not yet started, ongoing, or completed).

At this time, we recommend no direct manipulation of plant populations in Crab Lake. No aquatic invasive plant species are known to be present and no native plants exhibit nuisance population size or distribution.

Recommended Actions for the Crab Lake APM Plan

Action #1: Formally adopt the Aquatic Plant Management Plan.

Objective: To provide foundation for long-term native plant community conservation and stewardship and to be prepared for response to AIS introductions.

Monitoring: The Lake Association and Town Lakes Committee oversee activity and maintains the plan.

Status: Planned for 2015.

Action #2: Monitor water quality.

Objective: Continue with collection and analysis of water quality parameters to detect trends. Expand monitoring to include parameters for which little information exists (see Appendix C for individual parameters).

Monitoring: The Lake Association or Town Lakes Committee oversees activity and maintains data.

Status: Ongoing.

Action #3: Monitor the lake for aquatic invasive plant species.

Objective: To understand the lake's biotic community, provide for early detection of AIS and continue monitoring any existing populations of AIS.

Monitoring: The Lake Association or Town Lakes Committee oversees activity and maintains data.

Status: Ongoing.

Action #4: Monitor the lake for aquatic invasive animal species.

Objective: To understand the lake's biotic community, provide for early detection of AIS and continue monitoring any existing populations of AIS.

Monitoring: The Lake Association or Town Lakes Committee oversees activity and maintains data.

Status: Ongoing.

Action #5: Form an Aquatic Invasive Species Rapid Response Team and interface with the Town Lakes Committee AIS Rapid Response Coordinator.

Objective: To be prepared for AIS discovery and efficient response.

Monitoring: The Lake Association and/or Town Lakes Committee coordinate activity.

Status: Planned for 2015.

Recommended Actions for the Crab Lake APM Plan

Action #6: Conduct quantitative plant survey every five years using WDNR Point-Intercept Methodology.

Objective: To watch for changes in native species diversity, floristic quality, plant abundance, and plant distribution and to check for the occurrence of non-native, invasive plant species.

Monitoring: Town Lakes Committee (Wilderness Waters Program) oversees and maintains data; copies to WDNR.

Status: Anticipated in 2019.

Action #7: Update the APM plan approximately every five years or as needed to reflect new plant information from plant surveys and monitoring.

Objective: To have current information and management science included in the plan.

Monitoring: Lake Association and/or Town Lakes Committee (Wilderness Waters Program) oversees and maintains data; copies to WDNR.

Status: Ongoing.

Action #8: Develop a Citizen Lake Monitoring Network to monitor for invasive species and develop strategies including education and monitoring activities (see <http://www.uwsp.edu/cnr/uwexplakes/clmn> for additional ideas).

Objective: To create a trained volunteer corps to monitor aquatic invasive species and to educate recreational users regarding AIS.

Monitoring: The Lake Association oversees activity and reports instances of possible introductions of AIS.

Status: Anticipated to begin in 2015.

Action #9: Become familiar with and recognize the water quality and habitat values of ordinances and requirements on boating, septic, and property development.

Objective: To protect native aquatic plants, water quality, and riparian habitat.

Monitoring: Lake residents and other stakeholders.

Status: Ongoing.

Recommended Actions for the Crab Lake APM Plan

Action #10: Promote adherence to, and enforcement of, the Town of Presque Isle's 200 foot no-wake ordinances (from shoreline and islands).

Objective: To minimize recreational impacts on the aquatic plant community and shoreline habitats, and promote safe boating.

Monitoring: Town Lakes Committee oversees activity and assesses effectiveness.

Status: Ongoing.

Action #11: Create an education plan for the property owners and other stakeholders that will address issues concerning aquatic and riparian plant communities.

Objective: To educate stakeholders about issues and topics that affect the lake's aquatic and riparian plant communities, including topics such as: (1) the importance of the aquatic plant community; (2) no or minimal mechanical removal of plants along the shoreline is desirable and that any plant removal should conform to Wisconsin regulations; (3) the value of a natural shoreline in protecting the aquatic plant community and lake health; (4) nutrient sources to the lake and the role excess nutrients play in degradation of the aquatic plant community; (5) the importance of reducing or eliminating use of fertilizers on lake front property; (6) the importance of minimizing transfer of AIS to the lake by having dedicated watercraft and cleaning boats that visit the lake.

Monitoring: Town Lakes Committee oversee(s) activity and assesses effectiveness.

Status: Anticipated to begin in 2015.

Action #12: Monitor the lake watershed for purple loosestrife.

Objective: Identify purple loosestrife populations before they reach large size.

Monitoring: The Lake Association and/or Town Lakes Committee oversees activity.

Status: Anticipated in 2015.

Action #13: Monitor the cattail and *Phragmites* patches.

Objective: Confirm species identification and patches are increasing in area.

Monitoring: The Lake Association and/or Town Lakes Committee oversees activity with help from a consulting specialist as needed.

Status: Anticipated in 2015.

CHAPTER 6

Contingency Plan for AIS

Unfortunately, sources of aquatic invasive plants and other AIS are numerous in Wisconsin. Some infested lakes are quite close to Presque Isle Township. There is an increasing likelihood of accidental introduction of AIS to Presque Isle Township Lakes through conveyance of life stages by boats, trailers, and other vectors. It is important for the Town Lakes Committee and other lake stewards to be prepared for the contingency of aquatic invasive plant species colonization in a Presque Isle Township water body.

For riparian owners and users of a lake ecosystem, the discovery of AIS invokes a sense of tragedy that elicits an immediate desire to “fix the problem.” Although strong emotions may be evoked by such a discovery, a deliberate and systematic approach is required to appropriately and effectively address the situation. An aquatic plant management plan (one including a contingency plan for AIS) is the best tool by which the process can be navigated. In fact the APM plan is a requirement in Wisconsin for some kinds of aquatic plant management actions. One of the actions outlined in the previous chapter was to establish an Aquatic Invasive Species Rapid Response Team. This team and its coordinator are integral to the management process. It is important for this team to be multi-dimensional (or at least have quick access to the expertise that may be required). AIS invade not just a single lake, but an entire region since the new infestation is an outpost from which the AIS can more easily colonize other nearby water bodies. For this reason it is strategic for the Rapid Response Team to include representation from regional stakeholders.

Exhibit 10 provides a flowchart outlining an appropriate rapid response to the suspected discovery of an aquatic invasive plant species. The response will be most efficient if an AIS Rapid Response Team has already been established and is familiar with the contingency plan. In the remainder of this chapter we further describe the approach.

When a suspect aquatic invasive plant species is found, either the original observer or a member of the Rapid Response Team (likely the coordinator) should collect an entire plant specimen including roots, stems, and flowers (if present). The sample should be placed in a sealable bag with a small amount of water to keep it moist. Place a label in the bag written in pencil with date, time, collector’s name, lake name, location, town, and county. Attach a lake


map to the bag that has the location of the suspect AIS marked and GPS coordinates recorded (if GPS is available). The sample should be placed on ice in a cooler or in a refrigerator. Deliver the sample to the WDNR Water Resource Management Specialist (Kevin Gauthier in Woodruff), the Vilas County AIS Coordinator (Cathy Higley) or the WDNR Lakes Coordinator as soon as possible (at least within three days). The WDNR or their botanical expert(s) will determine the species and confirm whether or not it is an aquatic invasive plant species.

If the suspect specimen is determined to be an invasive plant species, the next step is to determine the extent and density of the population since the management response will vary accordingly. The Rapid Response Team should conduct (or have its consultant conduct) a survey to define the colony's perimeter and estimate density. If less than five acres (or <5% of the lake surface area), it is designated a "Pioneer Colony." If greater than five acres (or >5% of the lake surface area) then it is designated an "Established Population." Once the infestation is characterized, "at risk" areas should also be determined and marked on a map. For example, nearby boat landing sites and areas of high boat traffic should be indicated.

When "pioneer" or "established" status has been determined, the WDNR Lakes Coordinator should be consulted in order to determine appropriate notifications and management responses to the infestation. Determining whether hand-pulling or chemical treatment will be used is an important and early decision. Necessary notifications of landowners, governmental officials, and recreationists (at boat landings) will be determined. Whether the population's perimeter needs to be marked with buoys will be decided by the WDNR. Funding sources will be identified and consultants and contractors will be contacted where necessary. The WDNR will determine if a further baseline plant survey is required (depending on type of treatment). A post treatment monitoring plan will be discussed and established to determine the efficacy of the selected treatment.


Once the Rapid Response Team is organized, one of its first tasks is to develop a list of contacts and associated contact information (phone numbers and email addresses). At a minimum, this contact list should include: the Rapid Response Coordinator, members of the Rapid Response Team, County AIS Coordinator, WDNR Lakes Management Coordinator, Lake Association Presidents (or other points of contact), local WDNR warden, local government official(s), other experts, chemical treatment contractors, and consultant(s).

Exhibit 10. Aquatic Invasive Plant Species Rapid Response



Page left intentionally blank





Appendix A
Literature Cited

Page left intentionally blank



LITERATURE CITED

- Hauxwell, J., S. Knight, K. Wagner, A. Mikulyuk, M. Nault, M. Porzky and S. Chase. 2010. *Recommended baseline monitoring of aquatic plants in Wisconsin: sampling design, field and laboratory procedures, data entry and analysis, and applications*. Wisconsin Department of Natural Resources Bureau of Science Services, PUB-SS-1068 2010. Madison, WI.
- Natural Heritage Inventory. 2014. *Elements by TownRange for Vilas County*. Retrieved 2015. <http://dnr.wi.gov/topic/NHI/documents/Vilas_County.pdf>
- Nichols, Stanley A. 1999. *Floristic Quality Assessment of Wisconsin Lake Plant Communities with Example Applications*. *Journal of Lake and Reservoir Management* 15(2): 133-141.
- Premo, Dean, Angie Stine, Caitlin Clarke and Kent Premo. 2015. *The Wilderness Waters Adaptive Management Plan (Presque Isle Township, Vilas County, Wisconsin) - The Wilderness Waters Program*. White Water Associates, Inc.
- US Department of Agriculture, Natural Resources Conservation Service. June 1986. *Urban Hydrology for Small Watersheds*. Technical Release-55.
- Walters, C. 1986. *Objectives, constraints, and problem bounding*. In W.M. Getz, ed., *Adaptive Management of Renewable Resources*. Macmillan Publishing Company. New York, NY. p. 13+.
- Wisconsin Administrative Code NR-102-10 (1m). *Outstanding Resource Waters*. Revised 2010. Retrieved 2015. <http://docs.legis.wisconsin.gov/code/admin_code/nr/100/102.pdf#page=7>
- Wisconsin Administrative Code NR-01-05, 06, 07. Revised 2014. *Designated Waters*. Retrieved 2015. <https://docs.legis.wisconsin.gov/code/admin_code/nr/001/1/05>
- Wisconsin Department of Natural Resources. 2013. *Outstanding and Exceptional Resource Waters*. Retrieved 2015. <<http://dnr.wi.gov/topic/SurfaceWater/orwerw.html>>

Page left intentionally blank



Appendix B

Aquatic Plant Survey Tables and Figures

Table of Contents

Table 1. Summary statistics for point-intercept aquatic plant survey

Table 2. Plant species and distribution statistics

Figure 1. Number of plant species recorded at sample sites

Figure 2. Rake fullness ratings for sample sites

Figure 3. Maximum depth of plant colonization

Figure 4. Sampling sites less than or equal to maximum depth of rooted vegetation

Figure 5. Substrate encountered at point-intercept plant sampling sites

Figure 6. Aquatic plant occurrences for 2014 point-intercept survey data

Figure 7. Point-intercept plant sampling sites with emergent and floating aquatic plants

Figure 8-13. Distribution of plant species

Page left intentionally blank



Table 1. Summary statistics for the 2014 point-intercept aquatic plant surveys for Crab Lake.

Summary Statistic	Value	Notes
Total number of sites on grid	1204	Total number of sites on the original grid (not necessarily visited)
Total number of sites visited	564	Total number of sites where the boat stopped, even if much too deep to have plants.
Total number of sites with vegetation	58	Total number of sites where at least one plant was found
Total number of sites shallower than maximum depth of plants	239	Number of sites where depth was less than or equal to the maximum depth where plants were found. This value is used for Frequency of occurrence at sites shallower than maximum depth of plants.
Frequency of occurrence at sites shallower than maximum depth of plants	24.27	Number of times a species was seen divided by the total number of sites shallower than maximum depth of plants.
Simpson Diversity Index	0.92	A nonparametric estimator of community heterogeneity. It is based on Relative Frequency and thus is not sensitive to whether all sampled sites (including non-vegetated sites) are included. The closer the Simpson Diversity Index is to 1, the more diverse the community.
Maximum depth of plants (ft.)	10.50	The depth of the deepest site sampled at which vegetation was present.
Number of sites sampled with rake on rope	26	
Number of sites sampled with rake on pole	287	
Average number of all species per site (shallower than max depth)	0.47	
Average number of all species per site (vegetated sites only)	1.93	
Average number of native species per site (shallower than max depth)	0.47	Total number of species collected. Does not include visual sightings.
Average number of native species per site (vegetated sites only)	1.93	Total number of species collected including visual sightings.
Species Richness	29	
Species Richness (including visuals)	35	
Floristic Quality Index (FQI)	37.2	

Table 2. Plant species recorded and distribution statistics for the 2014 Crab Lake aquatic plant survey¹.

Common name	Scientific name	Frequency of occurrence at sites less than or equal to maximum depth of plants	Frequency of occurrence within vegetated areas (%)	Relative Frequency (%)	Number of sites where species found	Number of sites where species found (including visuals)	Average Rake Fullness
Fern pondweed	<i>Potamogeton robbinsii</i>	9.21	37.93	19.64	22	22	1.27
Wild celery	<i>Vallisneria americana</i>	7.11	29.31	15.18	17	22	1.00
Narrow-leaved bur-reed	<i>Sparganium angustifolium</i>	2.93	12.07	6.25	7	10	1.00
Muskgrasses	<i>Chara</i> sp.	2.51	10.34	5.36	6	6	1.00
Large-leaf pondweed	<i>Potamogeton amplifolius</i>	2.51	10.34	5.36	6	22	1.00
Dwarf water-milfoil	<i>Myriophyllum tenellum</i>	1.67	6.90	3.57	4	5	1.00
White water lily	<i>Nymphaea odorata</i>	1.67	6.90	3.57	4	15	1.00
Berchtold's pondweed	<i>Potamogeton berchtoldii</i>	1.67	6.90	3.57	4	6	1.00
Softstem bulrush	<i>Schoenoplectus tabernaemontani</i>	1.67	6.90	3.57	4	9	1.00
Water horsetail	<i>Equisetum fluviatile</i>	1.26	5.17	2.68	3	16	1.00
Pickereelweed	<i>Pontederia cordata</i>	1.26	5.17	2.68	3	15	1.00
Spiral-fruited pondweed	<i>Potamogeton spirillus</i>	1.26	5.17	2.68	3	4	1.00
Large purple bladderwort	<i>Utricularia purpurea</i>	1.26	5.17	2.68	3	3	1.00
Water marigold	<i>Bidens beckii</i>	0.84	3.45	1.79	2	2	1.00
Watershield	<i>Brasenia schreberi</i>	0.84	3.45	1.79	2	12	1.00
Needle spikerush	<i>Eleocharis acicularis</i>	0.84	3.45	1.79	2	2	1.00
Northern manna grass	<i>Glyceria borealis</i>	0.84	3.45	1.79	2	4	1.00
Quillwort	<i>Isoetes</i> sp.	0.84	3.45	1.79	2	2	1.00
Brown-fruited rush	<i>Juncus pelocarpus</i> f. <i>submersus</i>	0.84	3.45	1.79	2	4	1.00
Southern naiad	<i>Najas guadalupensis</i>	0.84	3.45	1.79	2	2	1.00
Variable pondweed	<i>Potamogeton gramineus</i>	0.84	3.45	1.79	2	6	1.00
Floating-leaf pondweed	<i>Potamogeton natans</i>	0.84	3.45	1.79	2	3	1.50
Arrowhead	<i>Sagittaria</i> sp.	0.84	3.45	1.79	2	5	1.00
Creeping spikerush	<i>Eleocharis palustris</i>	0.42	1.72	0.89	1	20	1.00

Frequency of occurrence within vegetated areas (%): Number of times a species was seen in a vegetated area divided by the total number of vegetated sites.

¹ Specimens of each species were collected, pressed and identification was confirmed by Dr. Robert Freckmann, University of Wisconsin-Stevens Point, on 9/11/2014.

Table 2. Continued.

Common name	Scientific name	Frequency of occurrence at sites less than or equal to maximum depth of plants	Frequency of occurrence within vegetated areas (%)	Relative Frequency (%)	Number of sites where species found	Number of sites where species found (including visuals)	Average Rake Fullness
Common waterweed	<i>Elodea canadensis</i>	0.42	1.72	0.89	1	2	1.00
Alternate-flowered water-milfoil	<i>Myriophyllum alterniflorum</i>	0.42	1.72	0.89	1	1	1.00
Nitella	<i>Nitella</i> sp.	0.42	1.72	0.89	1	1	1.00
Water smartweed	<i>Persicaria amphibia</i>	0.42	1.72	0.89	1	2	1.00
Common bladderwort	<i>Utricularia vulgaris</i>	0.42	1.72	0.89	1	1	1.00
Three-way sedge	<i>Dulichium arundinaceum</i>				Visual	6	
Spatterdock	<i>Nuphar variegata</i>				Visual	5	
Ribbon-leaf pondweed	<i>Potamogeton epihydrus</i>				Visual	3	
Pipewort	<i>Eriocaulon aquaticum</i>				Visual	1	
Water lobelia	<i>Lobelia dortmanna</i>				Visual	1	
Knotsheath sedge	<i>Carex retrorsa</i>				Visual	1	
Cattail	<i>Typha</i> sp.				Boat survey		
Northwest Territory sedge	<i>Carex utriculata</i>				Boat survey		
Blue flag iris	<i>Iris versicolor</i>				Boat survey		
Smooth sawgrass	<i>Cladium merisoides</i>				Boat survey		
Common reed	<i>Phragmites</i> sp.				Boat survey		

Frequency of occurrence within vegetated areas (%): Number of times a species was seen in a vegetated area divided by the total number of vegetated sites.

Figure 1. Number of plant species recorded at Crab Lake sample sites (2014).



Figure 2. Rake fullness ratings for Crab Lake sample sites (2014).



Figure 3. Maximum Depth of Plant Colonization in Crab Lake, 2014.




Figure 4. Crab Lake sampling sites less than or equal to maximum depth of rooted vegetation (2014).




Figure 5. Crab Lake substrate encountered at point-intercept plant sampling sites (2014).




Figure 6. Crab Lake aquatic plant occurrences for 2014 point-intercept survey data.




Figure 7. Crab Lake point-intercept plant sampling sites with emergent and floating aquatic plants (2014).



Figure 8. Distribution of plant species, Crab Lake (2014).




Figure 9. Distribution of plant species, Crab Lake (2014).




Figure 10. Distribution of plant species, Crab Lake (2014).




Figure 11. Distribution of plant species, Crab Lake (2014).




Figure 12. Distribution of plant species, Crab Lake (2014).





Figure 13. Distribution of plant species, Crab Lake (2014).



Appendix C

Review of Lake Water Quality

Table of Contents

Introduction.....	1
Comparison of Crab Lake with other datasets	1
Temperature	1
Dissolved Oxygen.....	4
Water Clarity.....	7
Turbidity	8
Water Color.....	9
Water Level.....	9
User Perceptions	10
Chlorophyll <i>a</i>	11
Phosphorus.....	12
Trophic State.....	13
Nitrogen	15
Chloride	15
Sulfate	15
Conductivity.....	16
pH.....	16
Alkalinity	17
Hardness.....	17
Calcium and Magnesium Hardness	17
Sodium and Potassium.....	17
Dissolved Organic Carbon.....	17
Silica	18
Aluminum	18
Iron.....	18
Manganese	18
Sediment	18
Total Suspended Solids.....	19
Aquatic Invasive Species	19
Resources	22

Page left intentionally blank



Review of Crab Lake Water Quality

Prepared by Angie Stine, B.S., and Caitlin Clarke, B.S., White Water Associates, Inc.


Introduction

Crab Lake is located in Vilas County, Wisconsin. It is a 909 acre drainage lake with a maximum depth of 60 feet. The Waterbody Identification Code (WBIC) is 2953500. The purpose of this study is to develop baseline data. Our goal is to collect existing water quality data to give us a starting point, and continue to monitor Crab Lake for a comparison of environmental and human changes. Water quality data was retrieved from the Wisconsin DNR SWIMS database between 1992 and 2014. On November 13, 2012 UW Stevens Point monitored various water quality parameters on Crab Lake. Secchi disk measurements were collected by Citizen Lake Monitoring Network (CLMN) volunteers from 1992 to 2014. Chlorophyll *a* and total phosphorus were also collected by CLMN volunteers from 1993 to 2014.

Comparison of Crab Lake with other datasets

Lillie and Mason's *Limnological Characteristics of Wisconsin Lakes* (1983) is a great source to compare lakes within our region to a subset of lakes that have been sampled in Wisconsin. Wisconsin is divided into five regions of sampling lakes. Vilas County lakes are in the Northeast Region (Figure 1) and were among 243 lakes randomly selected and analyzed for water quality.

Figure 1. Wisconsin regions in terms of water quality.



Temperature

Measuring the temperature of a lake at different depths will determine the influence it has on the physical, biological, and chemical aspects of the lake. Lake water temperature influences the rate of decomposition, nutrient recycling, lake stratification, and dissolved oxygen (D.O.) concentration. Temperature can also

affect the distribution of fish species throughout a lake. Figure 2 displays the water temperatures from April to May, 1998, 2003, 2007, 2008, 2011, and 2013. In May, 1998, 2003, and 2013 the temperatures began to stratify from surface to bottom. In June, Crab Lake began to stratify at 18 to 20 feet (Figure 3). In July, Crab Lake stratified between 18 and 25 feet (Figure 4). In August, Crab Lake stratified between 20 and 30 feet (Figure 5). Crab Lake water temperatures started mixing in September, and by October the temperature was consistent from surface to bottom (Figure 6). For more information about the influence of lake stratification on a lake's water quality and biota, please refer to *Understanding Lake Data* (Shaw et al., 2004).





Figure 4. Crab Lake temperature profile, July.




Figure 5. Crab Lake temperature profile, August.





Figure 6. Crab Lake temperature profile, September-October.



Dissolved Oxygen

The dissolved oxygen (D.O.) content of lake water is vital in determining presence of fish species and other aquatic organisms. Dissolved oxygen also has a strong influence on the chemical and physical conditions of a lake. The amount of dissolved oxygen is dependent on the water temperature, atmospheric pressure, and biological activity. Oxygen levels are increased by aquatic plant photosynthesis, but reduced by respiration of plants, decomposer organisms, fish, and invertebrates. The amount of dissolved oxygen available in a lake, particularly in the deeper parts of a lake, is critical to overall health.

In April and May, dissolved oxygen levels were high from surface water to bottom (Figure 7). In June, there was a notable variation of dissolved oxygen content (Figure 8). The June, 1995 and 1996 D.O. levels were lower than usual. In July, D.O. levels dropped between 20 and 30 feet (Figure 9). In August, D.O. levels decreased around 20 feet (Figure 10). In September and October, mixing occurred in the lake (Figure 11).

Figure 7. Crab Lake dissolved oxygen profile, April-May (2003, 2009-2011, 2013).




Figure 8. Crab Lake dissolved oxygen profile, June (1995-2013).




Figure 9. Crab Lake dissolved oxygen profile, July (1995-2009, 2011, 2013).




Figure 10. Crab Lake dissolved oxygen profile, August (1995-2013).





Figure 11. Crab Lake dissolved oxygen profile, September and October (1995-2007).




Water Clarity

Water clarity has two main components: turbidity (suspended materials such as algae and silt) and true color (materials dissolved in the water) (Shaw et al., 2004). Water clarity gives an indication of the overall water quality in a lake. Water clarity is typically measured using a Secchi disk (black and white disk) that is lowered into the water column on a tether. In simple terms, the depth at which the disk is no longer visible is recorded as the Secchi depth.

Figure 12 shows the July and August mean Secchi depths from 1992 to 2013. The shallowest mean Secchi depth was 8.5 feet in 2013, and the deepest reading was at 17 feet in 2007 (Figure 13). According to Table 1, Crab Lake’s 2013 mean Secchi depth is considered “fair,” with respect to water clarity.

Figure 12. Crab Lake Secchi depth averages (July and August only).



Past secchi averages in feet (July and August only).

(WDNR, 2014c)

Figure 13. Crab Lake’s July and August Secchi Data: Mean, Min, Max, and Secchi Count (1992-2013).

Year	Secchi Mean	Secchi Min	Secchi Max	Secchi Count
1992	14.1	14	14.5	5
1993	14.25	14	15	4
1994	15.38	14.5	17	4
1995	13.5	13	14	2
1996	13	13	13	2
1997	12.5	11	14	2
1998	13	13	13	1
1999	12.83	11	14	3
2000	13.5	13	14	2
2001	14.5	14	15	2
2002	13	12	14	3
2004	13.5	13	14	2
2005	15.5	15	16	2
2006	13.75	13	14.5	2
2007	17	17	17	1
2008	14	14	14	2
2009	15.5	15.5	15.5	1
2010	12	12	12	1
2011	10.13	9	12	4
2013	8.5	8	9	2

(WDNR, 2014c)

Table 1. Water clarity index (Shaw et al., 2004).


Water clarity	Secchi depth (ft.)
Very poor	3
Poor	5
Fair	7
Good	10
Very good	20
Excellent	32

Turbidity

Turbidity is another measure of water clarity, but is caused by suspended particulate matter rather than dissolved organic compounds (Shaw et al., 2004). Particles suspended in the water dissipate light and reduce the depth at which the light can penetrate. This affects the depth at which plants can grow. Turbidity also affects the aesthetic quality of water. Water that runs off the watershed into a lake can increase turbidity by introducing suspended materials. Turbidity caused by algae is the most common reason for low Secchi readings (Shaw et al., 2004). In terms of biological health of a lake ecosystem, measurements less than 10 Nephelometric Turbidity Units (NTU) represent healthy conditions for fish and other organisms. Crab Lake turbidity was analyzed in November, 2012 and had a value of 1 NTU.

While checking Secchi depth, the CLMN volunteers also rate water clarity and describe it as “clear” or “murky.” From 1997 to 2013, 92% of volunteers rated the water in Crab Lake as “clear” (Figure 14).

Figure 14. Crab Lake water appearance, 1997-2013.



Water Color

Color of lake water is related to the type and amount of dissolved organic chemicals. Its main significance is aesthetics, although it may also influence light penetration and in turn affect aquatic plant and algal growth. Many lakes have naturally occurring color compounds from decomposition of plant material in the watershed (Shaw et al., 2004). Units of color are determined from the platinum-cobalt scale and are therefore recorded as Pt-Co units. Shaw states that a water color between 0 and 40 Pt-Co units is low. Crab Lake water color was analyzed in November, 2012 and had a value of 14.3 Pt-Co.

CLMN volunteers also recorded their perceptions of water color in Crab Lake. Since 1992, 100% of volunteers indicated the water appeared “brown” in color.

Water Level

CLMN volunteers also recorded their opinion of the lake level as “high,” “normal,” or “low.” Figure 15 displays the CLMN opinions of water level from 1992-2013. In 1994, 1995, 1997 and 2006, 100% of volunteers viewed the water level as “low.” In 1996, 100% viewed the water level as “high.” In 2012, 100% of volunteers viewed the water level as “normal.”

In 2010, CLMN volunteers monitored Crab Lake water levels from May to September. Two water level records were also collected in 2013 (Figure 16).

Figure 15. Crab Lake water level percentage viewed by CLMN, (1992-2013).






Figure 16. Crab Lake water level, 2010 and 2013.



User Perceptions

When Secchi depth readings are collected, the CLMN record their perceptions of the water, based on the physical appearance and the recreational suitability. These perceptions can be compared to water quality parameters to see how the lake user would experience the lake at that time. When interpreting the transparency data, we see that when the Secchi depth decreases, the rating of the lake’s physical appearance also decreases. These perceptions of recreational suitability are displayed by year in Figure 17. From 1992 to 2007 and from 2009 to 2010, 100% of CLMN volunteers recorded Crab Lake to be “beautiful, could not be nicer.” From 2011 to 2013, the majority of volunteers said Crab Lake had “very minor aesthetic problems.”


Figure 17. Crab Lake aesthetic value, (1992-2013).



Chlorophyll *a*

Chlorophyll *a* is the photosynthetic pigment that makes plants and algae green. Chlorophyll *a* in lake water is therefore an indicator of the amount of algae. Chlorophyll *a* concentrations greater than 10 µg/L are perceived as a mild algae bloom, while concentrations greater than 20 µg/L are perceived as a nuisance. Chlorophyll *a* was analyzed on a routine basis in Crab Lake. Figure 18 displays the chlorophyll *a* values from 1993 to 2014 during the growing season (April-May) and summer season (June-August). As we can see, chlorophyll *a* values were well below nuisance levels and well below the average levels for Wisconsin natural lakes.


Figure 18. Crab Lake Chlorophyll *a*, (1992-2014).



Phosphorus

In more than 80% of Wisconsin's lakes, phosphorus is the key nutrient affecting the amount of algae and plant growth. If phosphorus levels are high, excessive aquatic plant growth can occur.

Phosphorus originates from a variety of sources, many of which are related to human activities. Major sources include human and animal wastes, soil erosion, detergents, septic systems and runoff from farmland or lawns (Shaw et al., 2004). Phosphorus provokes complex reactions in lakes. An analysis of phosphorus often includes both soluble reactive phosphorus and total phosphorus. Soluble reactive phosphorus dissolves in the water and directly influences plant growth (Shaw et al., 2004). Its concentration varies in most lakes over short periods of time as plants take it up and release it. Total phosphorus is considered a better indicator of a lake's nutrient status than soluble reactive phosphorus because its levels remain more stable (Shaw et al., 2004). Total phosphorus includes soluble phosphorus and the phosphorus in plant and animal fragments suspended in lake water. Ideally, soluble reactive phosphorus concentrations should be 10 µg/L or less at spring turnover to prevent summer algae blooms (Shaw et al., 2004). A concentration of total phosphorus below 20 µg/L for lakes should be maintained to prevent nuisance algal blooms (Shaw et al., 2004). Crab Lake total phosphorus values were collected from 1993 to 2014 during the growing and summer seasons (Figure 19). Total phosphorus levels remained at or below the nuisance threshold. According to Figure 21, the average total phosphorus for Crab Lake (10.5) can be categorized as "very good."



Total phosphorus was also analyzed at 45 feet (considered the hypolimnion) (Figure 20). In the hypolimnion, total phosphorus values were above the threshold value. The 2014 total phosphorus (hypolimnion) classifies Crab Lake as "good."





Figure 21. Total phosphorus concentrations for Wisconsin’s natural lakes and impoundments (Shaw et al., 2004).




Trophic State

Trophic state is another indicator of water quality (Carlson, 1977). Lakes can be divided into three categories based on trophic state – oligotrophic, mesotrophic, and eutrophic. These categories reflect a lake’s nutrient and clarity levels (Shaw et al., 2004).

Trophic State Index (TSI) was calculated by the WDNR using Secchi measurements (1992-2014), chlorophyll *a* (1993-2014), and total phosphorus (1993-2014) collected by the CLMN (Figure 22). The 2014 average chlorophyll *a* TSI (44) and total phosphorus TSI (48) classifies Crab Lake as “mesotrophic” (Table 2).

Figure 22. Crab Lake Trophic State Index (1992-2014).

Trophic State Index Graph



Monitoring Station: Crab Lake - Deep Hole, Vilas County
 Past Summer (July-August) Trophic State Index (TSI) averages.

● = Secchi	■ = Chlorophyll	▲ = Total Phosphorus
TSI(Chl) = TSI(TP) = TSI(Sec)		
It is likely that algae dominate light attenuation.		
TSI(Chl) > TSI(Sec)		
Large particulates, such as Aphanizomenon flakes dominate		
TSI(TP) = TSI(Sec) > TSI(Chl)		
Non-algal particulate or color dominate light attenuation		
TSI(Sec) = TSI(Chl) >= TSI(TP)		
The algae biomass in your lake is limited by phosphorus		
TSI(TP) > TSI(Chl) = TSI(Sec)		
Zooplankton grazing, nitrogen, or some factor other than phosphorus is limiting algae biomass		

(WDNR, 2014c)

30-40	Oligotrophic: clear, deep water; possible oxygen depletion in lower depths; few aquatic plants or algal blooms; low in nutrients; large game fish usual fishery
40-50	Mesotrophic: moderately clear water; mixed fishery, esp. panfish; moderate aquatic plant growth and occasional algal blooms; may have low oxygen levels near bottom in summer
50-60	Mildly Eutrophic: decreased water clarity; anoxic near bottom; may have heavy algal bloom and plant growth; high in nutrients; shallow eutrophic lakes may have winterkill of fish; rough fish common
60-70	Eutrophic: dominated by blue-green algae; algae scums common; prolific aquatic plant growth; high nutrient levels; rough fish common; susceptible to oxygen depletion and winter fishkill
70-80	Hypereutrophic: heavy algal blooms through most of summer; dense aquatic plant growth; poor water clarity; high nutrient levels

(WDNR, 2014c)

Researchers use various methods to calculate the trophic state of lakes. Common characteristics used to make the determination are: total phosphorus (important for algae growth), chlorophyll *a* concentration (a measure of the amount of algae present), and Secchi disk readings (an indicator of water clarity) (Shaw et al., 2004) (Table 3).

Table 3. Trophic classification of Wisconsin Lakes based on chlorophyll *a*, water clarity measurements, and total phosphorus values (Shaw et al., 2004).

Trophic class	Total phosphorus µg/L	Chlorophyll <i>a</i> µg/L	Secchi Disk (ft.)
Oligotrophic	3	2	12
	10	5	8
Mesotrophic	18	8	6
	27	10	6
Eutrophic	30	11	5
	50	15	4

Nitrogen

Nitrogen is second only to phosphorus as an important nutrient for aquatic plant and algae growth (Shaw et al., 2004). Human activities on the landscape greatly influence the amount of nitrogen in a lake. Nitrogen may come from lawn fertilizer, septic systems near the lake, or from agricultural activities in the watershed. Nitrogen may enter a lake from surface runoff or groundwater sources.

Nitrogen exists in lakes in several forms. Crab Lake was analyzed for total Kjeldahl nitrogen (0.37 mg/L in November, 2012), ammonium (0.03 mg/L in November, 2012), and nitrate-nitrite (0.04 mg/L in November, 2012). Nitrogen is a major component of all organic (plant and animal) matter. Decomposing organic matter releases ammonia, which is converted to nitrate if oxygen is present (Shaw et al., 2004). All inorganic forms of nitrogen can be used by aquatic plants and algae (Shaw et al., 2004). If these inorganic forms of nitrogen exceed 0.3 mg/L (as N) in spring, there is sufficient nitrogen to support summer algae blooms (Shaw et al., 2004). Elevated concentrations of ammonium, nitrate, and nitrite, derived from human activities, can stimulate or enhance the development, maintenance and proliferation of primary producers (phytoplankton, benthic algae, macrophytes), contributing to the widespread phenomenon of the cultural (human-made) eutrophication of aquatic ecosystems (Camargo et al., 2007). The nutrient enrichment can cause important ecological effects on aquatic communities, since the overproduction of organic matter, and its subsequent decomposition, usually lead to low dissolved oxygen concentrations in bottom waters, and sediments of eutrophic and hypereutrophic aquatic ecosystems with low turnover rates (Camargo et al., 2007).

Chloride

The presence of chloride (Cl^-) where it does not occur naturally indicates possible water pollution (Shaw et al., 2004). Chloride does not affect plant and algae growth and is not toxic to aquatic organisms at most of the levels found in Wisconsin (Shaw et al., 2004). Crab Lake chloride level was < 0.2 mg/L in November, 2012. The chloride concentration was well below the generalized distribution gradient found in surface waters in Wisconsin.

Sulfate

Sulfate in lake water is primarily related to the types of minerals found in the watershed, and to acid rain (Shaw et al., 2004). Sulfate concentrations are noted to be less than 10 mg/L in Vilas County (Lillie and

Mason, 1983). University of Wisconsin-Steven's Point laboratory sampled Crab Lake in November, 2012. They calculated a sulfur-ICP (inductively coupled plasma) value of 27.31 mg/L.

Conductivity

Conductivity is a measure of the ability of water to conduct an electric current. Conductivity is reported in micromhos per centimeter ($\mu\text{mhos/cm}$) and is directly related to the total dissolved inorganic chemicals in the water. Usually, values are approximately two times the water hardness, unless the water is receiving high concentrations of human-induced contaminants (Shaw et al., 2004). Crab Lake conductivity levels were sampled in November, 2012 with a conductance of 90 $\mu\text{mhos/cm}$.

pH

The acidity level of a lake's water regulates the solubility of many minerals. A pH level of 7 is considered neutral. The pH level in Wisconsin lakes ranges from 4.5 in acid, bog lakes to 8.4 in marl lakes (Shaw et al., 2004). Natural rainfall in Wisconsin averages a pH of 5.6. Some minerals become available under low pH (especially aluminum, zinc, and mercury) and can inhibit fish reproduction and/or survival. Mercury and aluminum are not only toxic to many kinds of wildlife, but also to humans (especially those that eat tainted fish). The pH scale is logarithmic, so every 1.0 unit change in pH increases the acidity tenfold. Water with a pH of 6 is 10 times more acidic than water with pH of 7. A lake's pH level is important for the release of potentially harmful substances and affects plant growth, fish reproduction and survival. A lake with neutral or slightly alkaline pH is a good lake for fish and plant survival. Crab Lake pH has not been tested, and should be considered in future water quality sampling.

Table 4 indicates the effects pH levels less than 6.5 will have on fish. While moderately low pH does not usually harm fish, the metals that become soluble under low pH can be important. In low pH waters, aluminum, zinc, and mercury concentrations increase if they are present in lake sediment or watershed solids (Shaw et al., 2004).

Table 4. Effects of acidity on fish species (Olszyk, 1980).

<i>Water pH</i>	<i>Effects</i>
6.5	Walleye spawning inhibited
5.8	Lake trout spawning inhibited
5.5	Smallmouth bass disappear
5.2	Walleye & lake trout disappear
5	Spawning inhibited in most fish
4.7	Northern pike, sucker, bullhead, pumpkinseed, sunfish & rock bass disappear
4.5	Perch spawning inhibited
3.5	Perch disappear
3	Toxic to all fish

Alkalinity

Alkalinity levels in a lake are affected by the soil minerals, bedrock type in the watershed, and frequency of contact between lake water and these materials (Shaw et al., 2004). Alkalinity is important in a lake to buffer the effects of acidification from the atmosphere. Acid rain has long been a problem with lakes that have low alkalinity levels and high potential sources of acid deposition. In November, 2012 alkalinity in Crab Lake was < 4 mg/L.

Hardness

Hardness levels in a lake are affected by the soil minerals, bedrock type in the watershed, and frequency of contact between lake water and these materials (Shaw et al., 2004). One method of evaluating hardness is to test for calcium carbonate (CaCO₃). Crab Lake hardness has not been tested, and should be included in future water quality sampling.

Calcium and Magnesium Hardness

The carbonate system provides acid buffering through two alkaline compounds: bicarbonate and carbonate. These compounds are usually found with two hardness ions: calcium and magnesium (Shaw et al., 2004). Calcium is the most abundant cation found in Wisconsin lakes. Its abundance is related to the presence of calcium-bearing minerals in the lake watershed (Shaw et al., 2004). Aquatic organisms such as native mussels use calcium in their shells. The aquatic invasive zebra mussel tends to need calcium levels greater than 20 mg/L to maintain shell growth. Crab Lake had a calcium level of 7.49 mg/L in November, 2012 which is an indication that zebra mussels could not flourish. Magnesium was also sampled in November, 2012 with a value of 2.25 mg/L.

Sodium and Potassium

Sodium and potassium are possible indicators of human pollution in a lake, since naturally occurring levels of these ions in soils and water are very low. Sodium is often associated with chloride and gets into lakes from road salting, fertilizations, and human and animal waste (Shaw et al., 2004). Potassium is the key component of commonly-used potash fertilizer, and is abundant in animal waste. Both of these elements are held by soils to a greater extent than is chloride or nitrate; therefore, they are not as useful as indicators of pollution impacts (Shaw et al., 2004). Although not normally toxic themselves, they provide a strong indication of possible contamination by more damaging compounds (Shaw et al., 2004). Crab Lake sodium (1.07 mg/L) and potassium (0.76 mg/L) were sampled in November, 2012.

Dissolved Organic Carbon

Dissolved Organic Carbon (DOC) is a food supplement, supporting growth of microorganisms, and plays an important role in global carbon cycle through the microbial loop (Kirchman et al., 1991). In general, organic carbon compounds are a result of decomposition processes from dead organic matter such as plants. When water contacts highly organic soils, these components can drain into rivers and lakes as DOC. DOC is also extremely important in the transport of metals in aquatic systems. Metals form extremely strong complexes with DOC, enhancing metal solubility while also reducing metal

bioavailability. Baseflow concentrations of DOC in undisturbed watersheds generally range from 1 to 20 mg/L carbon. Crab Lake DOC was analyzed in August 2014 (8.06 mg/L) and June 2014 (8.78 mg/L).

Silica

The earth's crust is abundant with silicates or other compounds of silicon. The water in lakes dissolves the silica and pH can be a key factor in regulating the amount of silica that is dissolved. Silica concentrations are usually within the range of 5 to 25 mg/L. Generally lakes that are fed by groundwater have higher levels of silica. Crab Lake silica levels have not been tested, and should be included in future water quality sampling.

Aluminum

Aluminum occurs naturally in soils and sediments. In low pH (acidic) environments aluminum solubility increases greatly. With a low pH and increased aluminum values, fish health can become impaired. This can have impacts on the entire food web. Aluminum also plays an important role in phosphorus cycling in lakes. When aluminum precipitates with phosphorus in lake sediments, the phosphorus will not dissolve back into the water column as readily. Crab Lake aluminum levels have not been tested, and should be included in future water quality sampling.

Iron

Iron also forms sediment particles that store phosphorus when dissolved oxygen is present. When oxygen concentration gets low (for example, in winter or in the deep water near sediments) the iron and phosphorus dissolve in water. This phosphorus is available for algal blooms. Crab Lake iron levels have not been tested, and should be included in future water quality sampling.

Manganese

Manganese is a mineral that occurs naturally in rocks and soil. In lakes, manganese is usually in particulate form. When the dissolved oxygen levels decrease, manganese can convert from an insoluble form to soluble ions. A manganese concentration of 0.05 mg/L can cause color and staining problems. Manganese data is unknown for Crab Lake, so future water quality sampling should include this parameter.

Sediment

Lake bottom sediments are sometimes analyzed for chemical constituents that they contain. This is especially true for potentially toxic metals such as mercury, chromium, selenium, and others. Lake sediments also tend to record past events as particulates settle down and become part of the sediment. Biological clues for the historic conditions in the lake can be gleaned from sediment samples. Examples include analysis of pollen or diatoms that might help understand past climate or trophic states in the lake. Sediment data was not collected for Crab Lake, and future sampling should include this parameter.

Total Suspended Solids

Total suspended solids are all particles suspended in lake water. Silt, plankton, and wastes are examples of these solids and can come from runoff of agricultural land, erosion, and can be produced by bottom-feeding fish. As the suspended solid levels increase, they absorb heat from sunlight which can increase the water temperature. They can also block the sunlight that plants need for photosynthesis. These events can in turn affect the amount of dissolved oxygen in the lake. Lakes with total suspended solids levels less than 20 mg/L are considered “clear,” while levels between 40 and 80 mg/L are “cloudy.” Total suspended solids data is unknown for Crab Lake, so future water quality sampling should include this parameter.

Aquatic Invasive Species

There are two invasive species found in Crab Lake: common reed and rusty crayfish.

Common reed, often referred to by its scientific name, *Phragmites*, is a perennial wetland grass that can grow 3-20 feet tall (WDNR, 2014b). It has stiff, hollow stems and roots that can reach up to 6 feet deep. It is generally found in disturbed areas and quickly becomes established by its spreading root system. It can form monocultures that push out native plant species, and can alter the hydrology, wildlife habitat and increase the potential for fires (WDNR, 2014b). *Phragmites* was discovered in Crab Lake in 2010. It should be noted that during the 2014 aquatic plant survey, specimens of *Phragmites* spp. were collected and were identified as native. More specimens within these *Phragmites* stands should be collected and identified.

Rusty crayfish are native to parts of Ohio, Tennessee, Kentucky and Indiana, and were likely introduced to Wisconsin waters by fishermen using the crayfish as bait (Gunderson, 2014). Rusty crayfish negatively affect other native crayfish species, cause destruction to aquatic plant beds, reduce fish populations by eating eggs, and cause shoreland owners recreational problems (Gunderson, 2014). It is illegal to possess both live crayfish and angling equipment simultaneously on any inland Wisconsin water (except Mississippi River) (WDNR, 2015). It is also illegal to release crayfish into a water body without a permit (WDNR, 2015). Rusty crayfish have been found in Averill Lake (2003), Big Lake (1961), McCullough Lake (2003), Papoose Lake (2003), Presque Isle Lake (2003), Rice Creek, Round Lake (1998), Stateline Lake (2011) and Van Vliet Lake (2003). Rusty crayfish were found in Crab Lake in 2009.

The University of Wisconsin-Madison’s Aquatic Invasive Species Smart Prevention program classifies Crab Lake as “Not Suitable” for zebra mussels, based on calcium and conductivity levels found in the lake (UW-Madison).

AIS monitoring has been conducted by a Citizen Aquatic Invasive volunteer in 2009 and 2010. In 2009, curly-leaf pondweed, Eurasian water-milfoil, zebra mussels, rusty crayfish and purple loosestrife were monitored. The volunteer said the rusty crayfish and purple loosestrife were present. No coordinates were recorded for the purple loosestrife observation, so future searches of purple loosestrife around Crab Lake would be beneficial. In 2010, the volunteer looked for some additional invasives such as the Chinese mystery snail, banded mystery snail, freshwater jellyfish, hydrilla, and the spiny waterflea. The only AIS found was the rusty crayfish.

Additional AIS monitoring was conducted in July and August, 2010 by the DNR using a zooplankton tow for zebra mussel veligers, with no finds. In September, 2012 an expansion for the AIS survey included a plankton tow for spiny and the fishhook water fleas, with no finds. The DNR also searched for purple loosestrife, *Phragmites*, flowering rush, hydrilla, Brazilian waterweed, Eurasian water-milfoil, curly-leaf pondweed, yellow floating heart, quagga mussels, zebra mussels, New Zealand mudsnail, faucet snail, red swamp crayfish, Chinese mystery snail, and banded mystery snail. Only *Phragmites* was present.

Clean Boats, Clean Waters (CBCW) is a program that inspects boats for aquatic invasive species and in the process educates the public on how to help stop the spread of these species. Clean Boats, Clean Waters inspected 321 boats on Crab Lake in 2014 with 513 people contacted (Figure 23). Some boaters were contacted more than once by CBCW and 100% were willing to answer questions (Figure 24). Ninety percent of patrons asked said they were on a different waterbody in the last five days. This high percentage makes education on how to stop the spread of AIS an important part of the CBCW. Figure 25 indicates that the majority of people understood the information. It would be beneficial to continue with the CBCW boat inspections to educate boaters about aquatic invasive species and to minimize the transfer of aquatic invasive species to other water bodies.

Figure 23. Crab Lake Clean Boats Clean Waters data (WDNR, 2014a).




Figure 24. Crab Lake Clean Boats Clean Waters data (WDNR, 2014a).





Figure 25. Crab Lake Clean Boats Clean Waters data (WDNR, 2014a).



Resources

- Camargo, Julio A., Álvaro Alonso (Lead Author); Raphael D. Sagarin (Topic Editor). 02 April 2007. *Inorganic nitrogen pollution in aquatic ecosystems: causes and consequences*. In: Encyclopedia of Earth. Eds. Cutler J. Cleveland (Washington, D.C.: Environmental Information Coalition, National Council for Science and the Environment). Retrieved 2015.
<http://www.eoearth.org/article/Inorganic_nitrogen_pollution_in_aquatic_ecosystems:_causes_and_consequences>
- Carlson, R.E. 1977. *A Trophic State Index for Lakes*. *Limnology and Oceanography*. 22: 361-369.
- Gunderson, Jeff. 2014. *Rusty Crayfish: A Nasty Invader*. Minnesota Sea Grant. Retrieved 2015.
<http://www.seagrant.umn.edu/ais/rustycrayfish_invader>
- Kirchman, David L.; Suzuki, Yoshimi, Garside, Christopher, Ducklow, Hugh W. (15). 1991. *High turnover rates of dissolved organic carbon during a spring phytoplankton bloom*. *Nature* 352 (6336): 612–614. Doi:10.1038/352612a0. Retrieved 2015.
<<http://www.nature.com/nature/journal/v352/n6336/abs/352612a0.html>>
- Lillie, R. A. and J. W. Mason. 1983. *Limnological Characteristics of Wisconsin Lakes*. Wis. Dept. of Natural Resources Tech. Bull. Page 138. Madison, WI.
- Olszyk, D. 1980. *Biological Effects of Acid Rain*. Testimony, Wis. Public Service Commission Docket No. 05-EP-2. 5 pp.
- Shaw, B. Mechenich, C, and Klessig, L. 2004. *Understanding Lake Data (G3582)*. Board of Regents of the University of Wisconsin System. Madison, WI.
- University of Wisconsin-Madison, Center for Limnology, Vander Zanden Lab. *Aquatic Invasive Species Smart Prevention*. Retrieved 2015. <<http://www.aissmartprevention.wisc.edu/>>
- Wisconsin Department of Natural Resources. 2014a. *Clean Boats, Clean Waters*. Retrieved 2015.
<<http://dnr.wi.gov/lakes/cbcw/>>
- Wisconsin Department of Natural Resources. 2014b. *Phragmites (Phragmites australis)*. Retrieved 2015.
<<http://dnr.wi.gov/topic/Invasives/fact/Phragmites.html>>
- Wisconsin Department of Natural Resources. 2014c. *Surface Water Integrated Monitoring Systems (SWIMS) Database*. Retrieved 2015. <<http://dnr.wi.gov/topic/surfacewater/swims/>>
- Wisconsin Department of Natural Resources. 2015. *Rusty Crayfish-Round Lake*. Retrieved 2015.
<<http://dnr.wi.gov/lakes/invasives/AISDetail.aspx?roiseq=66778265>>