



**\*For lakes/sites not snorkeled, substitute:**

Boat landing site - 15 rake throws and 15 D-net samples

Targeted site - 5 rake throws and 5 D-net samples

50 meander sites - 10 rake throws and 10 D-net samples during meander survey between sampling sites for a total of 50 meander survey sites

**\*\*If lake/site was not snorkeled, indicate why? (circle one) stained water, turbid water, blue-green bloom, chemical treatment, other \_\_\_\_\_.**

**\*\*\* Density Ratings**

- 1 - A few plants or invertebrates
- 2 - One or a few plant beds or colonies of invertebrates
- 3 - Many small beds or scattered plants or colonies of invertebrates
- 4 - Dense plant, snail or mussel growth in a whole bay or portion of the lake
- 5 - Dense plant, snail or mussel growth covering most shallow areas

**Step 2:** Collect Waterflea Tows from 3 sites: the deep hole (DH) and 2 other sites in water deeper than 15 feet (if possible).

Site	Depth sampled	Method (horizontal or oblique)	Net diameter (30 or 50 cm)	Ethanol added (Y or N)	Samples combined (Y or N)
7	4 ft	horizontal	50 cm		

→ 30m ropeline only

**Step 3:** Collect Veiliger Tows from 3 sites: the deep hole (DH), outlet site (OS), and or downwind site (DS) in water deeper than 4 meters (if possible).

Site	Depth sampled	Net diameter (30 or 50 cm)	Ethanol added (Y or N)	Samples combined (Y or N)
7	47	0.5m		

**Step 4:** Were voucher specimens submitted? Yes No (circle) If yes, where? (circle) Freckmann Herbarium, DNR Science Services, UW La Crosse, or Other \_\_\_\_\_

**Step 4:** Data was entered into SWIMS on 9/15/13 by Jennifer Stelwagen

**Step 5:** Data was proofed on 9/25/13 by Ryan Kottiff

**Notes:**